



Ricardo
Energy & Environment



Gibraltar City Inventory 2017

A City-Level Greenhouse Gas Emissions Inventory for Gibraltar

Report for HM Government of Gibraltar

Customer:

Sera Fromow, Department of the Environment,
HM Government of Gibraltar

Customer reference:

ED61636

Confidentiality, copyright & reproduction:

This report is the Copyright of the Government of Gibraltar. It has been prepared by Ricardo Energy & Environment, a trading name of Ricardo-AEA Ltd, under contract to the Government of Gibraltar dated 17/03/2016/17/03/2016/17/03/2016/17/03/2016.

The contents of this report may not be reproduced in whole or in part, nor passed to any organisation or person without the specific prior written permission of the Government of Gibraltar. Ricardo Energy & Environment accepts no liability whatsoever to any third party for any loss or damage arising from any interpretation or use of the information contained in this report, or reliance on any views expressed therein.

Contact:

Ellie Kilroy
Ricardo Energy & Environment
Gemini Building, Harwell, Didcot, OX11 0QR,
United Kingdom

t: +44 (0) 1235 75 3706

e: ellie.kilroy@ricardo.com

Ricardo-AEA Ltd is certificated to ISO9001 and ISO14001

Author:

Ellie Kilroy, Luke Jones, Rose Bailey, Charles Walker

Date:

17 May 2019

Ricardo Energy & Environment reference:

Ref: ED61636- Issue Number 1

Executive summary

Cities and communities present a significant opportunity in the management of global greenhouse gas (GHG) emissions; this has been increasingly recognised internationally and locally as organisations and initiatives increasingly champion city and community action, recognising that these places are often a focus of energy and resource consumption, create significant demands for mobility, and generate large quantities of waste. Globally, the focus is increasingly shifting towards enhancing the accounting and management of emissions at the city scale, and scaling up efforts to accurately monitor, report and verify activities as the basis for developing robust and evidence-based plans for action.

Since the landmark Paris Agreement in 2015, the emphasis is moving from making promises to taking action and tracking that action. Effective and committed governance at the national level will be key to achieving the Paris Agreement, however it is at the sub-national level where real gains in climate change mitigation will be made and communities like Gibraltar, with significant autonomy in key areas, have significant potential for leadership in demonstrating local level climate action.

The first step in managing Greenhouse Gas (GHG) emissions effectively at the city (or community) scale and making informed decisions to contribute to global mitigation efforts, is to have a good understanding of these emissions – the major sources, activities and relative contributions of different activities. Previous emissions data captured and reported through the UK Greenhouse Gas Inventory (GHGI) for Gibraltar did not give a complete picture of Gibraltar's GHG impact and mitigation activities, as its primary purpose is to monitor and report emissions produced on a territorial basis and is compiled following the Intergovernmental Panel on Climate Change (IPCC) 2006 guidelines to meet the UK's reporting commitments under the United Nations Framework Convention on Climate Change (UNFCCC). As it does not take account of activities outside of the geographic boundary or disaggregate information to a spatial resolution that is useful locally to enable targeted policy action such as electricity end use, the Gibraltar Emissions Inventory Programme was established to compile a detailed bottom-up inventory of community GHG emissions for Gibraltar annually from 2015.

This report covers the most recent inventory year, 2017. It considers emissions from all sources, including stationary combustion (both power generation and end consumption by sub-sector); mobile combustion (by road, marine, and shipping); waste disposal and wastewater; and industrial process and product use (IPPU) emissions. It follows the internationally approved standard from the Greenhouse Gas Protocol: the Global Protocol for Community-Scale Greenhouse Gas Emission Inventories (GPC) (Greenhouse Gas Protocol, 2014), and is reported using internationally approved tools.


Emissions are calculated for the seven Kyoto GHGs, reported as carbon dioxide equivalent (CO₂e) and are categorised by 'scope', to distinguish where emissions physically occur:

- Scope 1 emissions are directly emitted in boundary (direct emissions)
- Scope 2 emissions are indirectly emitted from in-boundary consumption of electricity (Indirect emissions)
- Scope 3 emissions are indirect and out of boundary emissions (Other direct emissions)

There are various levels of reporting, and this inventory also distinguishes between these different accounting approaches. The GPC has two reporting levels, known as BASIC and BASIC+, the latter including a greater number of sources, in particular some Scope 3 emission sources. It is recommended that cities aim to report BASIC+ emissions. Transboundary transport emissions are included under BASIC+ reporting, and this includes water-borne navigation. However, in the case of Gibraltar, much of this is international shipping (non-bunkering), and is excluded from the BASIC+ results presented in this report due to its very large impact on overall totals, and the lack of potential local influence. This sub-set can therefore be considered **Gibraltar's 'manageable emissions'**. This is shown in **Figure i**.

Sources that are deemed to be 'outside of scopes' (i.e. they are reported for information, but are not deemed to be within the influence or responsibility of Gibraltar – such as bunker fuel) would dominate emissions overall if included in emission totals, with bunker fuels accounting for 84.4% of total emissions when all are combined.

Figure i: BASIC+ “Manageable” emissions attributable to Gibraltar (under the GPC’s BASIC+ reporting, excluding transboundary international shipping)



The results for BASIC+ excluding transboundary (international) shipping present a picture much more aligned to those expected for a community, with stationary energy dominating, accounting for 53% of emissions. This is particularly the case in Gibraltar given that electricity generation is still predominantly produced using higher-carbon (diesel) fuel. Transportation also contributes just over one third of emissions, with 19% from road transport sources, 13% from aviation, and 4.3% from local boats. Waste and IPPU are smaller, at 6.7% and 3.6% respectively.

Scope 2 indirect emissions from electricity consumption are the largest source of emissions in Gibraltar, due to the reliance on electricity for all energy needs and generation technology. Because gas oil is used to generate electricity, the emissions per kilowatt hour (kWh) are considerably higher than, for example, the UK. The implied emission factor based on fuel consumption in power stations and total output is 0.69kg CO₂e/kWh in Gibraltar in 2017, compared with the UK 2017 grid factor of 0.35 kg CO₂e/kWh¹.

When comparing emissions with other global cities, per capita BASIC-level emissions are used (excluding any scope 3 emissions). For Gibraltar this equates to 6.3 tonnes CO₂e per person. This compares to a UK average of 5.5 tonnes per person in 2017, with the lowest cities emitting 3.1-3.4 tonnes per person².

When compiling the inventory for the latest year for Gibraltar, any improvements in data, methods or understanding are assessed and, where appropriate, are also applied to previous years inventories to enhance accuracy and consistency across the time series. The 2015 and 2016 inventories have therefore been revised, referred to as ‘2015r’ and ‘2016r’. More details on the revisions are found in the main report.

The main findings from the 2017 inventory are summarised as follows:

- ↓ Gibraltar’s total manageable emissions have decreased by 9% since 2015r and 8% since 2016r.

¹ <https://www.gov.uk/government/publications/greenhouse-gas-reporting-conversion-factors-2017>

² <http://www.centreforcities.org/reader/cities-outlook-2018/city-monitor/17-total-co2-emissions-per-capita/>

- ↓ Emissions from electricity generation have decreased by 19% since 2015r and 15% since 2016r; this is due to less fuel being used to generate a unit of electricity, implying improvements in efficiency at Gibraltar's electricity power stations. Electricity consumption by residents and activity in Gibraltar remained fairly consistent between 2015 and 2017.
- ↓ Emissions from Waste are around 13% and 10% lower in 2017 than 2015r and 2016r (respectively) due to a reduction in total waste arisings sent to landfill (and composting).
- ↓ Emissions from IPPU have decreased by 4% between 2015r and 2017; this follows trends in UK data that is used as a proxy for Gibraltar's emissions from product use (e.g. air conditioning and refrigeration).
- ↑ Emissions from road transport in Gibraltar have increased by 5% since 2015 due to more fuel being consumed by vehicles in Gibraltar.
- ↑ Emissions from scope 3 waterborne navigation* are 44% higher in the 2016r inventory than the 2015r inventory, largely due to longer liquid bulk container departing journey lengths in 2016; the reason for this is currently unknown. Trends in other vessel categories were more stable between the two years. These emissions have declined slightly since 2016r.
- ↑ Emissions from aviation are around 36% higher in 2017 than 2015r, likely due to an increased number of flights to London Gatwick, London Heathrow and Manchester between 2015r and 2016r. Between 2016r and 2017, aviation emissions have increased by 4%, largely due to an increased number of flights to London Gatwick and Manchester.

An emissions inventory is an ongoing tool for understanding and reporting emissions and allows the identification of major sources and priority areas for mitigation. It can be seen from the results presented above that there are some areas where mitigation efforts could be focussed in order to reduce greenhouse gas emissions, and some areas that are already demonstrating a reduction.

Stationary energy, as the highest contributor to overall manageable emissions, should be given priority and it is a positive sign that total emissions have declined 15% in a single year. In particular, the commercial and institutional sub-sector, as the highest end user category, could be further prioritised. Road transport makes up 19% of emissions, significant given the small size of the territory and the potential for policy interventions to reduce vehicle use. Finally, individual industries that are high-energy users, such as water supply, should be identified and ways to reduce energy consumption investigated. Shipping activity remains a large source of emissions; and one that will likely increasingly dominate the inventory even after excluding those emissions considered outside of scope.

This inventory has been compiled using the best available data and methods, however there remains potential for improvement, and future inventories should seek to build on the work undertaken here, and improve the accuracy, reliability, and coverage of data. This will better support decision making and targeted climate policy making, potentially bring co-benefits for other strategic areas through enhanced data capture and management on key activities, and enable the development of robust emission pathways modelling, supporting tracking of progress over time.

Table of contents

1	Introduction	7
1.1	Gibraltar’s climate commitments	8
2	The Global Protocol for Community-Scale Greenhouse Gas Emission Inventories (GPC)	9
2.1	Overview.....	9
2.2	Difference from national emissions reporting.....	9
2.3	Scopes and Sources	10
2.4	Reporting levels.....	12
2.5	Accounting and reporting principles	13
2.6	Data quality and notation keys	14
2.7	Accuracy.....	15
2.8	New reporting framework	15
3	Assessment boundaries	17
3.1	Geographic boundary	17
3.2	Temporal scale	17
3.3	Greenhouse gases reported.....	18
3.4	Sources and scopes	19
4	Calculation methodologies by emission source	21
4.1	Stationary energy	21
4.1.1	Energy industries: electricity generation	21
4.1.1.1	Summary of methods	21
4.1.1.2	Raw data	22
4.1.1.3	Determining emissions.....	22
4.1.2	Allocating emissions based on electricity consumption	24
4.1.2.1	Overview	24
4.1.2.2	Raw data	25
4.1.2.3	Determining activity.....	25
4.1.2.4	Determining emissions.....	26
4.1.3	Other stationary fuel combustion	27
4.2	Transportation	27
4.2.1	Road Transport	28
4.2.1.1	Overview	28
4.2.1.2	Raw data	29
4.2.1.3	Determining activity.....	31
4.2.1.4	Determining emissions.....	32
4.2.2	Marine – private boats.....	33
4.2.3	Shipping.....	34
4.2.3.1	Overview	34
4.2.3.2	Raw data	35
4.2.3.3	Determining activity.....	37
4.2.3.4	Determining emissions.....	42
4.2.4	Aviation.....	42
4.2.4.1	Overview	42
4.2.4.2	Raw data	43
4.2.4.3	Determining activity.....	43
4.2.4.4	Determining emissions.....	44
4.3	Waste	45
4.3.1	Summary	45
4.3.2	Raw data	46
4.3.2.1	Municipal solid waste	46
4.3.2.2	Clinical waste	47
4.3.3	Determining activity	47
4.3.3.1	Composition of MSW	47
4.3.4	Determining emissions	48
4.3.4.1	Solid waste disposal.....	48

4.3.4.2	Biological treatment.....	49
4.3.4.3	Clinical waste incineration.....	50
4.3.5	Wastewater	50
4.4	Industrial Processes and Product Use (IPPU)	52
4.4.1	Summary	52
4.4.2	Separating IPPU GHG emissions and energy-related GHG emissions	52
4.4.3	Determining activity	52
4.4.4	Estimating emissions.....	53
4.5	Agriculture, Forestry, and Other Land Use (AFOLU)	53
5	Results.....	54
5.1	Summary	54
5.2	Total emissions for Gibraltar	60
5.3	Total manageable emissions.....	60
5.4	Comparison with past inventories	61
5.4.1	Summary of changes	63
5.5	Reducing emissions in Gibraltar.....	66
5.6	Inventory accuracy	66
6	Recommendations.....	70
6.1	Improving inventory compilation and future year reporting	70
6.1.1	Improved activity data collection and management	70
6.1.2	Recalculations and tracking emissions over time	71
6.2	QAQC and verification.....	72
6.2.1	Quality control	72
6.2.2	Quality assurance.....	73
6.2.3	Verification.....	73

Appendices

Appendix 1: Common Reporting Framework (CRF) for 2017

Appendix 2: Detailed reasons for changes between 2017 and previous inventories

Appendix 3: Recalculations

Appendix 4: Data recommendations

Appendix 5: Data collection template

1 Introduction

Cities and communities present a significant opportunity in the reduction of global greenhouse gas (GHG) emissions; this has been increasingly recognised internationally and locally as organisations and initiatives increasingly champion city and community action, recognising that these places are often a focus of energy and resource consumption, create significant demands for mobility, and generate large quantities of waste. Globally, the focus is shifting towards enhancing the accounting and management of emissions at the city/community scale, and scaling up efforts to accurately monitor, report and verify activities as the basis for developing robust and evidence-based plans for action.

Since the landmark Paris Agreement in 2015, the emphasis is moving from making promises to taking action and tracking that action. Effective and committed governance at the national level will be key to achieving the Paris Agreement; however, it is at the sub-national level where real gains in climate change mitigation will be made. The Intergovernmental Panel on Climate Change's (IPCC) Special Report on Global Warming of 1.5°C (SR1.5) identifies cities and urban areas as one of four critical global systems that can accelerate and upscale climate action. Communities like Gibraltar, with significant autonomy in key areas, have significant potential for leadership in demonstrating local level climate action.

The first step in managing greenhouse gas (GHG) emissions effectively at the city (or community) scale and making informed decisions to contribute to global mitigation efforts, is to have a good understanding of these emissions – the major sources, activities and relative contributions of different activities. However, until 2014, cities lacked a common methodology for GHG accounting at the city scale. To overcome this problem, the Global Protocol for Community-Scale Greenhouse Gas Emission Inventories (GPC) was launched. The GPC offers cities, communities and local governments a robust, transparent and globally-accepted framework to consistently identify, calculate and report on city GHGs. It is methodologically consistent with national territory-based approaches to emissions accounting, but also provides the flexibility to account for emissions in ways that more accurately reflect local circumstances.

The 2013 city-level GHG inventory for Gibraltar³, prepared by Ricardo, was consistent with the draft version of the GPC standard available at the time. The 2013 inventory quantified emissions from stationary combustion by end user (power generation and consumption), mobile combustion (by road, marine and shipping), waste disposal and recycling, water supply, industrial emissions and indirect emissions associated with Gibraltar's supply chain (for instance, imports of food, construction materials and other goods). Gibraltar was one of the first communities to report a fully compliant GPC inventory, and the 2013 Gibraltar inventory was used as a case-study of best-practice in the final publication of the GPC, in meetings and with the World Bank and UNHABITAT, and in work with a number of global mega-cities (including Rio de Janeiro, Amman and Buenos Aires) on best practice in city GHG inventories. Gibraltar remains one of the leaders in community GHG reporting, having also reported an inventory for 2015 and 2016. Gibraltar is therefore part of a fast-growing number of urban communities establishing processes to report such data regularly. It is significant that Gibraltar has committed to reporting this data on an annual basis, to maintain its position as a leader in this field.

A key part of following the GPC guidelines is to update the inventory on a regular basis, ideally annually, as it is intended to be a 'live' tool for reporting, mitigating and tracking GHG emissions. Previous inventories should also be revised in line with updated methodologies or available data, to ensure an ongoing process of improvement and consistency and accuracy across the time series. This report therefore provides an update to the 2016 inventory (reported in 2018) for the year 2017, and identifies a number of improvements where recalculations of the 2015 and 2016 inventories have also been undertaken. This will also allow Gibraltar to continue showing best practice in city GHG inventories, successfully take part and report under initiatives such as the Global Covenant of Mayors for Climate and Energy, formerly known as the Compact of Mayors (see **Section 2**), and understand progress towards goals and inform the Climate Change Act and other programmes. Furthermore, this inventory programme will provide the evidence base to enable the undertaking of projections of emissions under different scenarios, including business-as-usual, planned policies and more ambitious actions.

³ https://www.gibraltar.gov.gi/new/sites/default/files/HMGoG_Documents/20150301-A_City-Level_Greenhouse_Gas_Inventory_for_Gibraltar_2013.pdf

This report and the accompanying GHG inventory data is part of Gibraltar's Emissions Inventory Programme (GibEmit), which in turn is part of the wider Gibraltar Air Quality and Climate programme, managed and delivered by Ricardo Energy & Environment.

1.1 Gibraltar's climate commitments

In October 2015, Gibraltar became a signatory of the Compact of Mayors, a global coalition of mayors and city officials pledging to reduce local GHGs, enhance resilience to climate change and track their progress transparently. The Compact collects the significant climate action data that cities are already reporting in a single, transparent platform. It therefore represents the greatest opportunity to bring attention to, and quantify, city action. As of January 2017, the Compact of Mayors merged with the EU's Covenant of Mayors to create the Global Covenant of Mayors for Climate and Energy (GCoM). GCoM brings together the world's two primary initiatives of cities and local governments – to advance city-level transition to a low emission and climate resilient economy, and to demonstrate the global impact of local action. Gibraltar is now one of over 9,200 cities and local governments who have committed to GCoM.

Under GCoM, Gibraltar have committed to regularly reporting a GHG Inventory, assessing climate risks and vulnerabilities, defining ambitious climate mitigation, resilience and energy targets, and creating a full climate action plan outlining how targets will be delivered, as depicted in Figure 1-1.

Figure 1-1: Compact of Mayors commitment requirements



Source: https://data.bloomberglp.com/mayors/sites/14/2015/07/Compact-of-Mayors-Full-Guide_July2015.pdf

It is important to note that as part of the merger, a common standard for city and local government GHG emissions inventory reporting for the GCoM, known as the 'Common Reporting Framework'⁴ (CRF), has been developed. More information on the CRF is presented in **Section 2.8**.

⁴ <https://www.globalcovenantofmayors.org/our-initiatives/data4cities/common-global-reporting-framework/>

2 The Global Protocol for Community-Scale Greenhouse Gas Emission Inventories (GPC)

2.1 Overview

In the past, inventory methods that cities have used have varied in the inclusion of emission sources and GHGs, how emissions sources are defined and categorised, and how transboundary emissions are treated. This inconsistency made comparisons between cities difficult; raised questions around data quality; and limited the ability to aggregate local, subnational and national government GHG emissions data. It was recognised that, to allow for more credible reporting, meaningful benchmarking and aggregation of climate data, greater consistency in GHG accounting was required. As noted in the **Introduction**, the GPC was launched in 2014 to address these issues and to offer a globally accepted robust and clear framework that builds on existing methodologies for calculating and reporting city-scale GHG emissions

Gibraltar's community-scale GHG inventory has been compiled following the GPC guidelines.

2.2 Difference from national emissions reporting

The GPC differs from national reporting methodologies (as required for reporting to the United Nations Framework Convention on Climate Change (UNFCCC)) in several fundamental ways, which reflect the unique circumstances of cities. Although adhering to basic principles of good practice in inventory compilation and reporting, the sources and sectors, and their categorisation are quite different. City-level emission inventories are not primarily focused on emissions from within the geographic boundary, as in a national inventory, but with emissions attributable to activities within the city. Therefore, a city-level inventory includes emissions that occur geographically outside the city (such as out of boundary waste disposal and transboundary transport). The focus on emission 'responsibility' also means that activities occurring in or near the city that are not the responsibility of the city can be excluded to give a more accurate picture of the city's impact; this is particularly significant for Gibraltar. Accounting for emissions on a territorial basis led to reports in summer 2012, based on data from the US Energy Information Administration, claiming that Gibraltar had the highest per capita carbon footprint in the world⁵; this was largely due to the volumes of bunker fuel sold to large marine cargo vessels⁶ compared with a small population. This presents a distorted view of GHG emission sources under local control in Gibraltar. An alternative city 'activity-based' approach to measure and report community-scale GHG emissions was needed for Gibraltar; hence Gibraltar's 2013 and 2015 community-scale GHG inventories and the 2016 inventory presented in this report.


Figure 2-1 shows, in simple terms, the difference in accounting approaches.

This 'responsibility' is broadly identified by means of 'scopes'. Scopes 1 and 2 are those sources occurring as a result of activities within the city boundary by and within the city. Scope 3 sources are those occurring, usually outside of the city boundary, as a consequence of activities. There is more detail on this in **Section 2.3**.

⁵ www.theguardian.com/environment/2012/jul/16/gibraltar-carbon-emissions-distorted-table

⁶ Bunker fuels refer to the storage and sale of fuels – typically gas oil and fuel oil – at national boundaries, in this case the trade of shipping fuels at the Port of Gibraltar.

Figure 2-1: Comparison between territorial accounting approach and GPC



Source: GPC v2.0 p.22

For Gibraltar, several key sources of emissions fall into the 'outside of scopes' category for a city inventory. These would be reported in a national inventory. Following an 'activity-based' approach which accounts for emissions that Gibraltar is 'responsible' for means that those sources that fall 'outside of scope' can be reported as such, and therefore excluded from inventory totals as a source beyond the responsibility of the community. Such sources for Gibraltar include bunker fuel sales and exported fuel in vehicles. Although methodologically more challenging to estimate (see methodology details below), it is important to attempt to differentiate between fuel used locally and that immediately exported by the many vehicles that cross the border to take advantage of cheaper fuel prices in Gibraltar.






2.3 Scopes and Sources

The GPC classifies emissions into six main sectors:

- I. Stationary energy
- II. Transportation
- III. Waste
- IV. Industrial processes and product use (IPPU)
- V. Agriculture, forestry, and other land use (AFOLU)
- VI. Other Scope 3 - Any other emissions occurring outside the geographic boundary as a result of city activities.

Emissions from these sectors are then sub-divided into sub-sectors and may be further divided into sub-categories. **Table 2-1** lists the GPC sectors and sub-sectors.

Table 2-1: GPC sectors and sub-sectors

Sector	Sub-sector
I. Stationary energy 	I.1 Residential buildings
	I.2 Commercial and institutional buildings and facilities
	I.3 Manufacturing industries and construction
	I.4 Energy industries
	I.5 Agriculture, forestry, and fishing activities
	I.6 Non-specified sources
	I.7 Fugitive emissions from coal
	I.8 Fugitive emissions from oil and natural gas systems
II. Transportation 	II.1 On-road
	II.2 Railways
	II.3 Waterborne navigation
	II.4 Aviation
	II.5 Off-road
III. Waste 	III.1 Solid waste disposal
	III.2 Biological treatment of waste
	III.3 Incineration and open burning
	III.4 Wastewater treatment and storage
IV. IPPU 	IV. 1 Industrial processes
	IV.2 Product use
V. AFOLU 	V.1 Livestock
	V.2 Land
	V.3 Aggregate sources
VI. Other Scope 3	VI.1 Other scope 3

Source: GPC


Activities taking place within a city can generate GHG emissions that occur inside the city boundary as well as outside the city boundary. To distinguish between these, the GPC groups emissions into three categories based on where they occur: scope 1, scope 2 or scope 3 emissions (**Table 2-2**).

Table 2-2: GPC inventory scopes

Scope	Definition	Example
Scope 1	GHG emissions from sources located within the city boundary.	<ul style="list-style-type: none"> Fuel consumed within the city boundary Waste generated and disposed of within the boundary
Scope 2	GHG emissions occurring as a consequence of the use of grid-supplied electricity, heat, steam and/or cooling within the city boundary.	<ul style="list-style-type: none"> Industrial consumption of grid-supplied electricity Residential consumption of grid-supplied heat
Scope 3	All other GHG emissions that occur outside the city boundary as a result of activities taking place within the city boundary.	<ul style="list-style-type: none"> Waste generated in the city but disposed in a landfill outside of the city Transmission and distribution losses from grid-supplied electricity

Sources and scopes of a GPC inventory are summarised in **Figure 2-2**.

Figure 2-2: Sources and scopes of a city GHG inventory



Source: GPC

2.4 Reporting levels

The GPC offers cities two levels of reporting demonstrating different levels of completeness, known as BASIC (**Figure 2-3**) and BASIC+ (**Figure 2-4**). The BASIC level covers emission sources that occur in almost all cities (Stationary Energy, in-boundary Transportation, and emissions from in-boundary generated Waste, including waste disposed outside the boundary). The BASIC+ level has a more comprehensive coverage of emissions sources (BASIC sources plus IPPU, AFOLU, transboundary transportation, and energy transmission and distribution losses) and reflects more challenging data collection and calculation procedures.

Gibraltar is reporting a BASIC+ inventory.

Figure 2-3: GPC reporting level - BASIC sources





Figure 2-4: GPC reporting level – BASIC+ sources



2.5 Accounting and reporting principles

Any inventory should include quality assurance/quality control (QA/QC) activities. The five key principles enshrined in the Intergovernmental Panel on Climate Change (IPCC) reporting guidelines, **transparency, consistency, comparability, completeness** and **accuracy** (TCCCA) should be adhered to in compiling inventory data and reports. The GPC also has five principles, although 'comparability' has been replaced with **relevance** and a more city-specific definition (see **Box 2-1**).

Box 2-1: GPC Principles of inventory compilation

1. **Relevance:** The reported GHG emissions shall appropriately reflect emissions occurring as a result of activities and consumption patterns of the city. The inventory will also serve the decision-making needs of the city, taking into consideration relevant local, subnational, and national regulations. The principle of relevance applies when selecting data sources, and determining and prioritizing data collection improvements.
2. **Completeness:** Cities shall account for all required emissions sources within the inventory boundary. Any exclusion of emission sources shall be justified and clearly explained. Notation keys shall be used when an emission source is excluded, and/or not occurring.
3. **Consistency:** Emissions calculations shall be consistent in approach, boundary, and methodology. Using consistent methodologies for calculating GHG emissions enables meaningful documentation of emission changes over time, trend analysis, and comparisons between cities. Calculating emissions should follow the methodological approaches provided by the GPC. Any deviation from the preferred methodologies shall be disclosed and justified.
4. **Transparency:** Activity data, emission sources, emission factors, and accounting methodologies require adequate documentation and disclosure to enable verification. The information should be sufficient to allow individuals outside of the inventory process to use the same source data and derive the same results. All exclusions shall be clearly identified, disclosed and justified.
5. **Accuracy:** The calculation of GHG emissions shall not systematically overstate or understate actual GHG emissions. Accuracy should be sufficient enough to give decision makers and the public reasonable assurance of the integrity of the reported information. Uncertainties in the quantification process shall be reduced to the extent that it is possible and practical.

Source: Section 2.1 of the GPC

2.6 Data quality and notation keys

Data collection is an integral part of developing and updating a GHG inventory. Data will likely come from a variety of sources and will vary in quality, format and completeness. In many cases, data will also need to be adapted for the purposes of the assessment. The GPC and the IPCC recognise these challenges and set out good practice data collection principles.

Not all data will be perfect, and there will be gaps, assumptions and limitations with data that are available. To recognise, accommodate and report these limitations, the GPC requires the use of notation keys (see **Table 2-3**). The GPC also requires that when notation keys are used, an accompanying explanation to justify the use of the notation key is also provided; this is to increase transparency and completeness. When collecting emissions data, it is important to establish first whether a source exists, and then the data availability and quality.

- If the source does not exist, 'NO' is used to indicate it is 'not occurring'. For example, in Gibraltar, there is no rail transport and no agriculture.
- If the activity does occur in the city, and data are available, then the emissions should be estimated. However, if the data are also included in another emissions source category or cannot be disaggregated, the notation key 'IE' would be used to indicate 'included elsewhere' and avoid double counting. The category in which they are included should be identified. For example, in Gibraltar emissions from water are included under stationary energy as the only emissions attributable to water are from the consumption of electricity.
- If the data are not available and, therefore, the emissions are not estimated, the notation key 'NE' would be used to indicate 'not estimated'. This is only permitted for scope 3 sources and IPPU and AFOLU sectors, reported under BASIC+ and therefore considered 'optional'.

Table 2-3: Use of notation keys

Notation key	Definition	Explanation
NO	Not occurring	An activity or process does not occur or exist within the city.
IE	Included elsewhere	GHG emissions for this activity are estimated and presented in another category of the inventory. That category should be noted in the explanation.
NE	Not estimated	Emissions occur but have not been estimated or reported; justification for exclusion should be noted.
C	Confidential	GHG emissions which could lead to the disclosure of confidential information and can therefore not be reported.

Source: Table 2.1 of the GPC

The GPC also requires a qualitative assessment of data quality to be reported; this involves using expert judgement to assign a rating of high (H), medium (M) or low (L) quality to the both the activity data and emission factors used in emission calculations (see **Table 2-4**).

Table 2-4: Data quality assessment

Data Quality	Activity data	Emission factor
High (H)	Detailed activity data	Specific emission factors
Medium (M)	Modelled activity data using robust assumptions	More general emission factors
Low (L)	Highly-modelled or uncertain activity data	Default emission factors

Source: Table 5.3 of the GPC

2.7 Accuracy

Most major emission sources within the Gibraltar inventory ultimately fall under electricity consumption (relevant to most Stationary Energy sub-sectors) or fuel consumption (such as road and marine sub-sectors), for which accurate totals are available from the power stations and import statistics, respectively. Therefore, these data sources act as the high level 'fuel balance' that is allocated across different sources from available activity data. This ensures that there is a high level of reliability in the total emission figures and double-counting is avoided. Any uncertainty is then associated with the activity data and allocation methods across different end users. Accuracy here is important for policy purposes, but less important for understanding the total amount of GHGs emitted.

2.8 New reporting framework

As mentioned, GCoM have recently released a common standard for city and local government GHG emissions inventory reporting for the GCoM, known as the 'Common Reporting Framework'⁷ (CRF). The CRF ensures a common approach for cities to monitor their performance against their individual action plans and targets while simultaneously creating a mechanism to transparently track the contributions and impacts of cities and local governments within the framework of the Paris Agreement. It helps track the invaluable contributions of all subnational actors, allow for comparison between jurisdictions and increase the potential for financing opportunities at the local, regional, and global levels. This common standard has emerged from a need to harmonise the GPC reporting undertaken under the Compact of Mayors, and the alternative accounting approaches promoted by the Covenant of Mayors. Whilst very similar, there are some slight differences and GCoM has sought to harmonise these through the CRF. The principles and requirements of the CRF are largely the same as the GPC, with amendments confined to nomenclature of categories and mandatory versus optional requirements.

⁷ <https://www.globalcovenantofmayors.org/our-initiatives/data4cities/common-global-reporting-framework/>

As Gibraltar reports one of the most complete GHG inventories of any community, the changes in reporting do not require any additional effort with regards to inclusion of sources, gases etc., nor affect the reporting of the 'manageable' emissions as per this report.

The main differences between the GPC and CRF that affect how Gibraltar's emissions will be reported are outlined below. Please note, these changes affect how emissions are reported and not the emissions themselves.

- Under the CRF, emissions are reported as '**direct**' and '**indirect**' emissions to distinguish where they physically occur, rather than using scopes as in the GPC. Under the CRF, emissions are categorised as:
 - **Direct emissions** (GPC Scope 1) due to fuel combustion in the buildings, equipment/facilities and transportation sectors within the city boundary. These emissions physically occur inside the city boundary.
 - **Other direct emissions** (GPC Scope 3) that are not related to fuel combustion, including: fugitive emissions from disposal and treatment of waste (including wastewater) generated within the city boundary, which may occur inside or outside the city boundary, and; fugitive emissions from natural gas distribution systems (such as equipment or pipeline leaks).
 - **Indirect emissions** (GPC Scope 2) due to consumption of grid-supplied energy (electricity, heat or cold) within the geographic boundary. Depending on where energy is generated, these emissions may physically occur inside or outside the city boundary.
- **Energy Industries** (GPC Sub-sector I.4.4) has been split into types of generation to enable reporting of how electricity is generated and the type of facilities generating electricity
- **Non-specified sources** (GPC Sub-sector I.6) have been removed and emissions are to be reported in one of the other sub-sectors

For the purposes of this report and Gibraltar's city inventory programme, emissions are still reported using the GPC. Appendix 1 reports Gibraltar's emissions following the CRF, as reported to CDP as part of Gibraltar's GCoM commitment. As mentioned, changes to reporting from following the CRF only affect how emissions are reported, and not the emissions themselves.

3 Assessment boundaries

This section sets out the reporting boundaries and requirements of the inventory.

3.1 Geographic boundary

The inventory is defined geographically by the territorial boundary of Gibraltar, as shown in **Figure 3-1**.

Figure 3-1: Map of Gibraltar



Source: <http://www.geoportal.gov.gi/webviewer/>

3.2 Temporal scale

This inventory covers all atmospheric emissions during calendar year 2017. Where 2017 data were not available, the most recent year's data have been used and the timescale noted accordingly. In particular, these are:

- Population: 2015 figure extrapolated to 2017 following recent trend (2017 not yet available)
- Gross domestic product (GDP): 2016 figure extrapolated to 2017 following recent trend (2017 not yet available)

3.3 Greenhouse gases reported

As per the GPC, Gibraltar accounts for emissions of the seven gases currently required for most national GHG inventory reporting under the Kyoto Protocol: carbon dioxide (CO₂), methane (CH₄), nitrous oxide (N₂O), hydrofluorocarbons (HFCs), perfluorocarbons (PFCs), sulfur hexafluoride (SF₆), and nitrogen trifluoride (NF₃). Nitrogen trifluoride was not one of the six gases originally mandated under the Kyoto Protocol, but was added for the second compliance period (starting 2012). The gases required by the GPC are the same seven gases currently required for most national GHG inventory reporting. CO₂ from biogenic sources are reported separately and not included in inventory totals.

The International System of Units (SI units) is used for measuring and reporting activity data, and all GHG emissions data are reported as metric tonnes of CO₂ equivalents (CO₂e). CO₂e accounts for the global warming potential (GWP) when measuring and comparing GHG emissions from different gases. Individual GHGs are converted into CO₂e by multiplying by the 100-year GWP coefficients given in the 2006 Intergovernmental Panel on Climate Change (IPCC) Guidelines (see **Table 3-1**). These are taken from the IPCC.

Gibraltar is using the 4th Assessment Report GWP values, consistent with the UK national GHG inventory and international best practice.

Box 3-1: Biogenic CO₂

Biogenic emissions are those that result from the combustion of biomass materials that naturally sequester CO₂, including materials used to make biofuels (e.g. crops, vegetable oils, or animal fats). For the purposes of national level GHG inventories, land-use activities are recorded as both sinks and sources of CO₂ emissions. Reporting emissions from combusting these biogenic fuels would result in double counting on a national level. The GPC also records land-use changes, and combusted biofuels may be linked to land-use changes in its own inventory, or other cities' inventories.

Under the CRF, reporting biogenic CO₂ is now optional.

Source: Box 4.2 of the GPC

Table 3-1: Global warming potentials (GWP) used in calculations, adapted from IPCC 2006 Guidelines.

Industrial designation or common name	Chemical formula	Lifetime (years)	Radiative efficiency (W m ⁻² ppb ⁻¹)	Global warming potential for given time horizon (100 years)
Carbon dioxide	CO ₂		1.4x10 ⁻⁵	1
Methane	CH ₄	12	3.7x10 ⁻⁴	25
Nitrous oxide	N ₂ O	114	3.03x10 ⁻³	298

3.4 Sources and scopes

Table 3-2 indicates the sources included in the inventory under each emission scope, and **Figure 3-2** shows this in diagrammatic format.

Table 3-2: Sources included in the inventory under each emission scope

Scope	Definition
Scope 1 (direct emissions)	All GHG emissions from sources located within the boundary of the city: <ul style="list-style-type: none"> • Stationary fuel combustion. • Direct IPPU emissions⁸ • Electricity generation (information item only). • Mobile fuel combustion: <ul style="list-style-type: none"> – Road vehicles in-boundary. – Marine vessels in-boundary. – Aircraft in-boundary⁹. – Rail in-boundary⁹. • Waste disposal. • AFOLU⁹.
Scope 2 (indirect emissions)	All GHG emissions occurring as a consequence of the use of grid-supplied electricity , heating and/or cooling within the city boundary: <ul style="list-style-type: none"> • Industrial electricity consumption⁹. • Commercial/other non-domestic electricity consumption. • Electricity consumption for key users (for example, water¹⁰).
Scope 3 (Other direct emissions)	All other GHG emissions that occur outside the city boundary as a result of activities within the city's boundary: <ul style="list-style-type: none"> • Mobile fuel combustion: <ul style="list-style-type: none"> – Road vehicles transboundary. – Marine vessels transboundary. – Aircraft transboundary. – Rail transboundary⁹. • Waste disposal and wastewater treatment.
Outside of scopes (reported under 'VI. Other Scope 3') (Other direct emissions)	Sources that occur in or within the vicinity of Gibraltar, but which occur indirectly as a result of activities outside the control or influence of the community. These sources are reported as information items and not included in the overall emission total: <ul style="list-style-type: none"> • Export of road transport fuels. • Marine fuel bunkering.

⁸ Industrial Process emissions are not occurring in Gibraltar. Product Use emissions are reported however.

⁹ Not occurring in Gibraltar

¹⁰ In Gibraltar, water emissions are included under Scope 2 as emissions are solely those associated with electricity consumption for desalination plant and pumping. No mains water is imported.

Figure 3-2: Gibraltar's emission sources by scope



4 Calculation methodologies by emission source

4.1 Stationary energy

Stationary energy is a significant part of any inventory. This is generally divided into two categories – emissions from stationary combustion of fuel, and emissions from generation and consumption of grid supplied electricity. Stationary energy sources appear in all reporting scopes as shown in **Box 4-1**.

Box 4-1: Stationary energy sources

Scope 1: Emissions from in-boundary emissions from fuel combustion and fugitive emissions.

- Combustion of fuels in buildings and industry.
- Conversion of primary energy sources in refineries and power plants (including production of electricity used by the power plant).
- Fossil resource and exploration within the city boundary.
- Fugitive emissions from fuel systems.

Scope 2: Emissions from the consumption of grid-supplied electricity, steam, heating and cooling.

Scope 3: Other out-of-boundary emissions.

- Transmission and distribution losses of electricity
- Steam, heating and cooling (not occurring in Gibraltar).

4.1.1 Energy industries: electricity generation


Electricity is the major energy source for Gibraltar and is the only energy industry present. Gibraltar is self-sufficient in electricity and operates as a 'closed system' (that is, there are no imports or exports from neighbouring regions). This allows for a very accurate calculation of the electricity-related emissions for Gibraltar.

4.1.1.1 Summary of methods

Electricity production includes two categories, which should add up to total emissions from fuel combusted for energy generation:

1. Electricity generation sold and distributed: this comprises emissions from all fuel use for electricity generation from main activity producers.
2. Auxiliary energy use on the site of energy production facilities.

The process of estimating emissions from electricity generation is shown in **Figure 4-1**.

Figure 4-1: Process of estimating emissions from electricity generation

4.1.1.2 Raw data

Raw data were obtained from the Gibraltar Electricity Authority (GEA) and consisted of electricity output, fuel use, fuel type and time period of reporting.

Gibraltar's power stations are:

- Waterport power station;
- North Mole Turbines, or temporary generators;
- OESCO power station; and,
- GMES power station.

All power stations use gas oil (either marine or automotive gas oil). Emission factors for fuels are taken from the UK National Atmospheric Emissions Inventory (NAEI) (2017 data) and are shown in **Table 4-1**.

Table 4-1: Emission factors for power station gas oil fuel (from UK NAEI 2017)

Pollutant	Unit	Emission factor
Carbon	kt/Mt fuel consumed	870
Methane (CH ₄)	kt/Mt fuel consumed	0.13
Nitrous oxide (N ₂ O)	kt/Mt fuel consumed	0.026

4.1.1.3 Determining emissions

To calculate emissions from electricity generation, total annual fuel use at the power stations by type is summed and multiplied by the relevant emission factor for each pollutant; the UK NAEI emission factors for gas oil has been used (**Table 4-1**). This figure is then multiplied by the pollutant's global warming potential (GWP) (or 44/12 to convert from carbon to CO₂) to give total carbon dioxide equivalent (CO₂e) emissions in tonnes. This gives the total emissions from generation. As emissions here are calculated

from consumption of a known quantity and type of fuel, and not from other activity data, it is possible to aggregate emissions.

To disaggregate emissions on an end-user basis, based on electricity consumption, an implied emission factor (IEF) calculated from known activity data is required. To calculate the IEF, total emissions associated with the fuel consumed to produce the electricity is divided across the total production of electricity to estimate emissions per unit. This then gives an estimate of the emissions for each unit consumed, in kt CO₂e per gigawatt hour (GWh) (as shown in **Table 4-2**). This IEF can then be multiplied by total electricity consumed (billings data) to give emissions from energy consumed by end-users.

The difference between electricity produced by the power stations and the electricity supplied to the Gibraltar electricity network is assigned to use of their own power at the power station sites.

The difference between the amount of electricity supplied to the Gibraltar electricity network and the amount of electricity that is billed for by AquaGib is assumed to be the transmission and distribution losses across the network.

Table 4-2: Implied emission factors for Gibraltar's power generation for 2017

Pollutant	Unit	IEF	IEF factor (CO ₂ e)
Carbon	kt/GWh	0.18869	0.6919
CH ₄	kt/GWh	0.00003	0.0007
N ₂ O	kt/GWh	0.00001	0.0017
Total	kt/GWh		0.6942

Figure 4-2: Turbines at North Mole



Figure 4-3: New skid-mounted generators at North Mole (2014)




4.1.2 Allocating emissions based on electricity consumption

Allocation of emissions from electricity generation to the end user uses data on total electricity consumption in Gibraltar and the IEF calculated for generation as the basis for calculations. Gibraltar is unusual in that all electricity consumed is also generated within the boundary. Therefore, total emissions data are allocated across different sectors.

4.1.2.1 Overview

A summary of the process is illustrated in **Figure 4-4**.

Figure 4-4: Summary of the process of calculating emissions

4.1.2.2 Raw data

A number of data sources were used in compiling estimates of emissions from electricity consumption. These were:

- GWh billings by tariff supplied by AquaGib (see **Table 4-3**).
- Electricity consumption data for key sectors, including hotels, the hospital, the airport
- Proxy data on employment by sector from the 2017 Employment Survey Report¹¹ (see **Table 4-4**). This shows proportion of employees by sector and allocation of industries to tariffs.

4.1.2.3 Determining activity

Electricity consumption data need to be allocated to end users through known consumption or an allocation based on a proxy indicator. Known consumption for sectors include domestic (residential) consumers (from AquaGib tariff data); hotel billings data (obtained directly from hotels); hospital and airport consumption, Ministry of Defence consumption and AquaGib water electricity billings. Known billings were subtracted from total billings data.

Remaining billings data are then allocated to sectors based on employment numbers from the the 2017 Employment Survey Report, and this employment data was used as a multiplier to billings data within tariff categories as shown in **Table 4-4**.

As mentioned, transmission and distribution losses are assumed to be the difference between the electricity that is supplied and the electricity that is billed. This is allocated to GPC sub-sectors based on the share of billed electricity consumption of each respective sub-sector.

The difference between electricity produced by the power stations and the electricity supplied to the Gibraltar electricity network is assigned to use of their own power at the power station sites.

¹¹ https://www.gibraltar.gov.gi/new/sites/default/files/HMGoG_Documents/Employment%20Survey%20Report%202017.pdf

Table 4-3: AquaGib electricity tariffs

Tariff number	Tariff name	Description	GPC sub-sector allocation
1	Lighting	Tariff for public lighting only	I.2.2
2	Power	Tariff for power only – examples include temporary sockets	I.2.2
3	Domestic	Residential properties only	I.1.2
4	Commercial	Majority of public sector and commercial premises (e.g. hospital)	I.2.2
5	Industrial maximum demand	Energy-intensive users, in particular bakeries, super markets, hotels	I.2.2
6A	Off-peak	Power during off-peak hours only	I.2.2
6B	Off-peak	Power during off-peak hours only	I.2.2
9	MOD Offices and Residential	Power used in MOD offices and residences	I.6.2

Table 4-4: Employment numbers by industrial/commercial sector, used as proxy data for electricity allocation

Industry	2017 employment	% of total
Shipbuilding	297	1%
Other Manufacture	226	1%
Electricity and Water Supply	277	1%
Construction	3472	12%
Wholesale and Retail Trade	3298	13%
Hotels and Restaurants	1763	7%
Transport and Communication	1549	6%
Financial Intermediation	2128	8%
Real Estate and Business Activities	3156	13%
Public Administration and Defence	2246	8%
Education	1080	4%
Health and Social Work	2375	9%
Other Services	4277	17%
Total	26144	100%

4.1.2.4 Determining emissions

Emissions are calculated by multiplying the GWh assigned to each end-user sector as above, by the IEF for each pollutant and its GWP, to give a value of CO₂e by end-user sector.

4.1.3 Other stationary fuel combustion

Scope 1 emissions from combustion of fuels in power stations in Gibraltar are covered above.

There is believed to be a small amount of stationary fuel combustion, in the form of bottled gas, assumed to be used in restaurants, hotels and the hospital. Fuel import data provided by HM Customs in 2015 has been used to estimate emissions from this source. In the absence of new data for the 2016 and 2017 inventories, we have assumed that the same fuel consumption occurred for 2017 and 2016 as 2015; this is an appropriate assumption as significant annual trends for this source are not expected. The import statistics refer to 'Petroleum gases and others gaseous hydrocarbons'; this is assumed to be LPG. Activity data is multiplied by the Defra Conversion Factors 2017 emission factor for LPG.

It is also understood that the hospital, airport and some hotels have fuel combustion capacity (such as diesel and gas oil used for back-up generators and LPG for cooking and patio heaters). New data was collected for the 2016 and 2017 inventories from a number of hotels and the hospital to reflect this activity and subtracted from the import statistics to avoid double counting. For the 2017 inventory, new data was also collected for the airport where fuel is consumed for back-up generation and for powering the aviation training simulator. When new data is identified, revisions are made to previous year's inventories to ensure completeness, assuming trends in usage have been unchanged.


There is no stationary fuel combustion in households as all energy requirements are met through electricity, so this source is not occurring (NO).

There is no fossil resource or exploration in Gibraltar, so this source is NO.

4.2 Transportation

The transport sector covers a wide range of emission sources, including road, rail (not present in Gibraltar), air and water, and consists of in-boundary and transboundary sources. In the case of Gibraltar, some transport sources (exported road and marine bunker fuels) are also estimated but excluded from totals.

Figure 4-5: End-use of imported fuel by sector, for petrol and automotive diesel assumed by the inventory



4.2.1 Road Transport

Road transport emissions have been calculated from Gibraltar's fuel import statistics for 2017. This effectively provides an 'energy balance' for total road transport fuel consumption. Imported fuel data is provided by the Port Authority of Gibraltar and is reallocated to different road vehicle types through a series of assumptions, further discussed below.

A proportion of this imported fuel also goes to private marine use. The allocation of this is discussed in **Section 4.2.2**. Although there is uncertainty in allocation, the overall fuel total and, therefore, emissions, particularly for CO₂, remains accurate.

4.2.1.1 Overview


Figure 4-6 gives a brief overview of how estimates of emissions due to road transport have been made. A more detailed explanation is given in the following sections.

For the highest emitting sources, Gibraltar-specific data have been sought and used. In some cases, for less significant sources, emission factors have been taken from the UK inventory. Generally speaking, these assumptions are based on factors that are unlikely to vary much between Gibraltar and the UK or the impact of any significant differences would be small.

Road transport emissions from fuel used by Gibraltar vehicles are assigned to Scope 1.

Road transport emissions from fuel used by non-Gibraltar vehicles are assigned to outside of scopes.

There is no way of differentiating transboundary transport (Scope 3).

Figure 4-6: Road transport method summary

4.2.1.2 Raw data

The licensing statistics provided by the Government of Gibraltar give a number of key pieces of information, allowing the nature of the road transport situation in Gibraltar to be determined. Particular data used were:

- The type of vehicle:
 - This allowed a decision on what kind of vehicle the record corresponded to and, in some cases, allowed a decision to be made about the fuel or weight class.
- Registration date:
 - This helped determine when vehicles were likely to have been manufactured and, hence, what European emission standard they will have been required to meet.
- The fuel type (that is, petrol or diesel vehicles).
- Cylinder capacity:
 - This was used to help determine the weight classes of the vehicles.
- The model and make:
 - Used to spot-check some assumptions and to correct other details (such as vehicle type) when found to be inaccurate.

Licensing statistics for the 2017 fleet were not made available to the inventory agency, and as such, a fleet turnover model was developed to approximate the shift in fleet composition between 2017 and

2016. This enabled the modelling the penetration of Euro 6 vehicles and the removal of older, less fuel-efficient vehicles from Gibraltar's road fleet. The turnover rate for this model was estimated using vehicle-specific survival rates implied by the 2016 licensing statistics, and vehicle sales assumptions were based on information in the Abstract of Statistics publication series.

Fuel import data for 2017 provided a high-level total energy consumption to allocate by transport mode. Prior to use in this inventory, the fuel import statistics required cleaning since the recorded mass and volume often implied an infeasible fuel density, suggesting that inconsistent units were used by importers when recording this data. Further details on the allocation of fuel use to the road transport sector are found in **Section 4.2.1.3**.

Fuel consumption allocated to road transport in 2017 is shown in **Table 4-5**.

Table 4-5: Gibraltar total road transport fuel use for 2016

Year	Fuel	Thousands of litres
2017	Motor spirits	14,087
2017	Automotive Gas Oil	27,183

Surveys of fuel stations carried out by the Department of the Environment in 2013, and then later in 2017, provide a snapshot of fuel use by vehicle type (commercial, private (assumed car) and motorcycle), the fuel type, and whether the vehicle is registered to Gibraltar or elsewhere (most typically Spain). Results from these surveys are combined and averaged to generate an estimate of fleet composition. This is because the results between the two surveys differed significantly, far beyond the extent that might be expected at typical fleet turnover rates, and so interpolating results between 2013 and 2017 would be misleading and likely highly inaccurate. Instead, the use of the 2017 results in this year's inventory is considered an expansion of the sample size and therefore, its representativeness to Gibraltar's fleet population. Some key differences are shown in **Box 4-2** below.

In addition, given the updated forecourt survey and differences identified, the road transport emissions for the 2015 inventory were revised following this new methodology and assumptions.

Box 4-2: Key differences between fuel station forecourt survey years and assumptions made

Key differences between the forecourt surveys were:

- The 2017 survey does not appear to count motorbikes for the majority of the dataset. To overcome this, it was assumed that the same proportion of motorcycles filled up between 2014 and 2017 and assumed that all two-wheelers were recorded as private vehicles in 2017 to offset this.
- There appears to be major shifts between 2014 and 2017 in the proportion of private vehicles originating from Gibraltar and from outside Gibraltar. After the correction to motorcycles, the 2017 % of private petrol vehicles from outside Gibraltar drops from 32% to 4%. This seems unlikely and is likely a reflection of the small sample size.
- There is a concurrent increase in the % of private petrol vehicles from Gibraltar from 34% to 62%.
- Diesel commercial vehicles registered to Gibraltar increased from 12% to 24%.

There is no obvious reason why the fleet composition of Gibraltar would have shifted significantly between 2014 and 2017, so using an interpolated time-series would be inaccurate and misleading as it would suggest a genuine trend. Therefore, we have used an average of the two surveys, suggesting that we do not think the fleet composition will have changed between these years (and hence 2015 and 2016 inventories).

Data was also available for fuel consumption for 2017 for one major bus company based in Gibraltar. In the 2015 inventory, data was ascertained for several other transport modes, including Government of Gibraltar vehicles, customs vehicles and fuel consumption from a second major bus company. This data was not available for 2017 and so it has been assumed that fuel consumption has not differed between the two years for these modes.

4.2.1.3 Determining activity

Road transport emissions are most accurately estimated from fuel consumption when the carbon content, and thus CO₂ emitted when combusted, is accurately known (although other pollutants are more greatly affected by the method of combustion). There are reliable data from the fuel import statistics for this. However, for a local-scale inventory, an understanding of how these emissions are allocated across modes by activity is more useful for informing policy. In this inventory, the fuel import data have been allocated to the road transport sector by vehicle and fuel type.

It was assumed that fuel import data, by motor spirit (petrol) and automotive gas oil (diesel) for 2017, included fuel use, but excluded use in private marine vessels. This is because marina fuel is tax exempt and so would not be included in import records. In the 2018 inventory submission, it was assumed that diesel and petrol consumption in small marina craft, such as speedboats and dingy craft, equated to 30% and 10% of imports of these fuels respectively. However, after conversations with the Gibraltar Port Authority¹², these assumptions are felt to overestimate marina fuel use. As a result, at least until more comprehensive data is made available for the inventory, a revised assumption of 15% and 5% for diesel and petrol marina use is made.

Time-series analysis of import statistics to Gibraltar since 2003 suggest that marina consumption is additional to the total imports for road transport use. From 2008-2012 it is believed that import statistics include both road transport and marine uses and so estimates of total private vessel consumption can be calculated using the offset to the import trends over this period. Therefore, for the 2017 inventory we now approximate that in addition to the 15% assumption for diesel oil, the amount of petrol consumed by these smaller vessels equates to 5% of total imports.

Table 4-6 - Fuel import data scaled and allocated for marine and road use

Fuel type	Net fuel imports (thousands of litres)	Use by private marine vessels (thousands of litres)	Total use (thousands of litres)
Motor spirit	14,087	704	14,791
Automotive gas oil	27,183	4,077	31,260

Data was provided by one of Gibraltar's two principle bus companies which allowed for the estimation of fuel consumption directly from this vehicle type. 2015 data for the other of these companies is available and total fuel consumption from this company is not assumed to have changed. In this case, the company could only provide data on annual fuel costs and so assumptions were made on the price paid for fuel to estimate fuel consumption. In addition, CO₂ emissions from urea consumption in Euro 6 buses was included in the inventory for the first time.

In the absence of vehicle activity data (e.g. mileage by mode) to assign fuels to vehicle classes, vehicle fleet data were used to calculate activity data (vkm travelled) by category. Forecourt survey data was processed and normalised to give a frequency of vehicle type (shown in **Table 4-7**) and, therefore, a fuel-use split. These fuel-use splits were then applied to total fuel use by type (as above), to give fuel use in kt by vehicle type – Gibraltar and non-Gibraltar.

Due to the lack of specific activity data for journey types (in-boundary and transboundary, and Gibraltar and non-Gibraltar vehicles) all fuel use by Gibraltar vehicles was assigned as in-boundary. All fuel use by non-Gibraltar vehicles was assigned to out-of-boundary with no explicit transboundary proportion (although some of the in-boundary fuel may be transboundary by Gibaltarians crossing the frontier).

¹² Personal communication with John Ghio, Deputy Captain of Port, March 2019

Table 4-7: Average fleet composition as indicated by the 2013 and 2017 forecourt surveys

Fuel	Gibraltar/non-Gibraltarian	Vehicle type	Average fleet composition by fuel type (%)
Diesel	Gibraltarian	Private vehicle	29%
		Commercial vehicle	18%
		Motorcycle ¹³	0%
	Non-Gibraltarian	Private vehicle	48%
		Commercial vehicle	3%
		Motorcycle ¹³	2%
Petrol	Gibraltarian	Private vehicle	48%
		Commercial vehicle	8%
		Motorcycle	10%
	Non-Gibraltarian	Private vehicle	18%
		Commercial vehicle	1%
		Motorcycle	16%

4.2.1.4 Determining emissions

Carbon emissions factors are derived using COPERT fuel consumption factors and weighted using the detailed fleet composition information as suggested by active vehicles listed in Gibraltar's licensing statistics, based on vehicle type, fuel used, weight class, European emission standard and, if applicable, catalyst type. As discussed in **Section 4.2.1.2**, licensing statistics for 2017 were not available for the 2019 compilation cycle and so a fleet turnover model was developed to model the penetration of Euro 6 vehicles and removal of older, and typically less fuel-efficient vehicles. This model was developed using a turnover rate assumption implied by the 2016 licensing statistics and the number of vehicles sold and registered in Gibraltar based on information within the Abstract of Statistics publication series. It has been assumed that all traffic in Gibraltar in traffic is urban and that the fleet characteristics of the non-Gibraltarian fleet not captured by the licensing statistics does not differ significantly. The emission factors then derived from the fuel consumption factors are the same as those used in the UK NAEI road transport projection models (using carbon contents provided by the United Kingdom Petroleum Industry Association, UKPIA).

Emission factors for methane and nitrous oxide are also the same as those used in the UK NAEI road transport projection models and are derived from the Transport Research Laboratory (TRL) emission factors for fuel consumption. Emissions are then calculated for each pollutant by multiplying the implied vkm travelled (shown in **Table 4-9**) by the fleet weighted emission factors.

Emissions from non-Gibraltarian vehicles are accounted for under 'Other Scope 3' and are therefore not included in BASIC or BASIC+ inventory totals.

¹³ Diesel motorcycles are reallocated to petrol in the final calculations as they are considered rare and are probably errors in the survey results.

Table 4-8: Fleet-weighted emission factors

Vehicle type	Weighted emission factor (g/km)		
	CO ₂	CH ₄	N ₂ O
Petrol cars	180	0.034	0.013
Diesel cars	189	0.002	0.015
Petrol LGVs*	277	0.025	0.013
Diesel LGVs	223	0.003	0.015
HGV **	593	0.062	0.020
Bus	667	0.077	0.022
Motorcycles	77	0.073	0.002

*Light goods vehicle (LGV) ** Heavy goods vehicle (HGV)

Table 4-9: Calculated fleet-weighted fuel consumption and vkm

Gibraltar/n on-Gibraltarian	Vehicle type	Fuel type	Fuel consumption (g/km)	Total calculated fuel consumption (kt)	Implied mvkm* travelled
Gibraltarian	Private vehicle	Petrol	57.6	4.54	77.42
Gibraltarian	Commercial vehicle	Petrol	88.4	0.77	8.74
Gibraltarian	Motorcycle	Petrol	24.6	0.98	39.59
Gibraltarian	Private vehicle	Diesel	60.2	6.41	106.44
Gibraltarian	Commercial vehicle	Diesel	79.1	4.11	51.99
Gibraltarian	Bus	Diesel	212.9	0.38	1.79
Non-Gibraltarian	Private vehicle	Petrol	57.6	1.69	28.89
Non-Gibraltarian	Commercial vehicle	Petrol	88.4	0.06	0.71
Non-Gibraltarian	Motorcycle	Petrol	24.6	1.79	72.57
Non-Gibraltarian	Private vehicle	Diesel	60.2	10.66	177.14
Non-Gibraltarian	Commercial vehicle	Diesel	79.1	0.71	8.97

*million vehicle kilometres

4.2.2 Marine – private boats

As noted above, a proportion of gas oil and petrol included in the fuel import data is used by private boats. This has been estimated at 15% and 5% of total demand for gas oil and petrol respectively in 2017, based on historic fuel import statistics from 2008 to 2012. Emissions have been estimated using the emission factors for marine gas oil and petrol as used within the UK NAEI. As activity data in this sector are not available, there is no way of allocating to specific activities within the private marine sector.

Figure 4-7: There are a large number of private boats in Gibraltar, but no bottom-up activity data are available on their fuel use and the characteristics of the resident fleet



4.2.3 Shipping


Shipping generates a large proportion of Gibraltar's emissions in the national inventory because of the considerable amount of bunkering activity and the fact that Gibraltar is a large international port near a major shipping lane. In this inventory, shipping is divided into two main categories: bunkering, and non-bunkering (that is, ships that call at Gibraltar with a purpose other than just obtaining fuel).

Shipping emissions from non-bunkering traffic are assigned to scope 3.

Shipping emissions from bunkering traffic are assigned to outside of scopes

4.2.3.1 Overview

Figure 4-8 gives a brief overview of how estimates of emissions due to shipping have been made. A more detailed explanation is given below.

Figure 4-8 – Process of estimating emissions from shipping

4.2.3.2 Raw data

The raw dataset was provided by the Gibraltar Port Authority and provides information on the shipping movements of all vessels that 'interface' with Gibraltar, including details of ship-type, gross tonnage, last port, and next port destination. However, a number of vessels included within this dataset carry out 'off port limit' calls, and do not enter Gibraltar waters; these are excluded from the dataset on the basis of additional information provided by the Port Authority. The key pieces of information used in the subsequent inventory calculations are:

- A distance (km) travelled to/from Gibraltar.
 - This is calculated using <http://ports.com/sea-route> to estimate the distance in nautical miles and converted to km. A weighted, ship-type specific average distance is derived to estimate more representative vessel journey lengths. The activity for ships travelling both to and from Gibraltar has been calculated, but only one direction (departing) should be included in Gibraltar's emissions total as per the GPC methodology. The origin and destination are those reported on the ship manifests.
 - The method is weighted according to the frequency at which boats visit various ports (and also applied to the 2015 revised inventory). In addition, averages for each of the ship types considered is calculated separately so that more characteristic distances are calculated. This causes a reduction in implied average journey distance since the majority of boats leaving the port visit nearby ports and therefore onward journeys are significantly shorter than the average previously estimated.
- Ship class
 - The given ship type was assigned to one of the below groups of ship, allowing the use of Tables 3-4 and 3-7 in the EMEP/EEA air pollutant emission inventory guidebook 2016¹⁴ Section 1.A.3.d Navigation. (See **Table 4-10**) is within or outside the scope of the inventory
 - Guidance from the Port Authority was used to determine which ships should be included within the inventory, and which were involved with either bunkering, or off port limit calls. **Table 4-11** illustrates the allocation on the basis of the registered purpose of call within the dataset.

¹⁴ www.eea.europa.eu/publications/emep-eea-guidebook-2016

Table 4-10 – Ship classification based on the EMEP/EEA Guidebook 2016¹⁴

Ship types	
Liquid bulk ships	Dry bulk carriers
Container	General cargo
Ro Ro Cargo	Passenger
Fishing	Tug
Other	

Table 4-11 – Definition of in-scope and out-of-scope shipping activity on the basis of stated purpose of call

Purpose of call	
In-scope	Out-of-scope
To Supply Bunkers	Bunkers
Arrested	Hold Inspection
Repairs	Slops Discharge
STS With Mother Ship (Bunker Barges only)	Crew Change
Laid Up	Underwater Cleaning
Waiting Orders	Medical Assistance
Gibraltar/Tangiers Ferry	Spares
Owners Change	Stores
Cruise Call	Charts
Stationed	Lub-Oil
STS	Provisions
Containers Loading/Unloading	Surveyor/Technician Transfer
Cargo Loading/Unloading	Underwater Inspection
Yacht Delivery	Cargo Sampling
Rocks Unloading/Loading	Change of Schedule
MOD Movement	Class Survey
Yacht Loading/Unloading	Bunker Survey
Eastern Anchorage - Awaiting Berth/Supply	Debunkers
Ship Sanitation Certificates	Pratique Note
Vehicle Loading/Unloading	Water Receive
Sail Training Ship Visit	Port Clearance Note
Eastern Anchorage - Awaiting STS	Shelter
Publicity Event	Compass Adjusting
Cancelled operation	Deliver Fenders
Dredging Works	STS Equipment Return
Sea Trials	PSC Inspection
STS Aegean	Underwater Survey
Waste Discharge	Tender/Service

Purpose of call	
In-scope	Out-of-scope
	PSC Mandatory Expanded Inspection
	Yacht Visit
	Detention
	Towing
	Under Tow
	Garbage Discharge
	Load Line Certificate
	Fuel Discharge
	Mid-Harbour Marina Berthing
	Radio Repairs
	Gyro Repairs
	Fenders Discharge

4.2.3.3 Determining activity

The key activity data of interest are the mass of fuel used, as this is the activity for which emissions factors are available within the UK NAEI. After processing the raw data from the port statistics, the activity dataset is in km. To convert this to a fuel use, it is possible to use the following to calculate fuel use using Equation 4-1.

- Fuel use per unit energy given in Table 3-4 of the shipping chapter of the EMEP/EEA air pollutant emission inventory guidebook 2016 (shown here in **Table 4-12**).
- The engine type weightings provided in Table 3-7 of the guidebook (shown here in **Table 4-13**).
- The main engine power in table 3-6 of the guidebook (shown here in **Table 4-14**).
- The average speeds in table 3-14 of the guidebook (shown here in **Table 4-15**).

Bunker fuel sales data also was supplied. However, the data did not closely correlate to the reported fuel import/export data also provided and was considered to be less robust, so was not used in the shipping inventory calculations.

Equation 4-1: Calculation for shipping emissions

$$\text{Fuel use (t)} = \frac{\text{Distance travelled(km)} * \text{Power}^{15} \text{ (kW)} * \text{Fuel use factor(t/kWh)}}{\text{Average speed (km/hour)}}$$

¹⁵ Main engine power

Figure 4-9: Bunkering ships

Table 4-12: Tier 2 emissions factors for shipping from the EMEP/EEA air pollutant emission inventory guidebook 2016¹⁴ (appears as Table 3-4 in guidebook)

Tier 2 default emission factors							
Engine type	Fuel type	NO _x 2000 (kg/tonne)	NO _x 2005 (kg/tonne)	NO _x 2010 (kg/tonne)	TSP - PM ₁₀ (kg/tonne)	PM _{2.5} (kg/tonne)	Specific fuel consumption (g fuel/kWh)
Gas turbine	BFO	20.0	19.3	18.6	0.3	0.3	305
	MDO/MGO	19.7	19.0	18.3	0.0	0.0	290
High-speed diesel	BFO	59.6	57.7	55.6	3.8	3.4	213
	MDO/MGO	59.1	57.1	55.1	1.5	1.3	203
Medium-speed diesel	BFO	65.7	63.4	61.3	3.8	3.4	213
	MDO/MGO	65.0	63.1	60.6	1.5	1.3	203
Slow-speed diesel	BFO	92.8	89.7	86.5	8.7	7.8	195
	MDO/MGO	91.9	88.6	86.5	1.6	1.5	185
Steam turbine	BFO	6.9	6.6	6.4	2.6	2.4	305
	MDO/MGO	6.9	6.6	6.4	1.0	0.9	290

Source: Entec (2002), Entec (2007), emission factors calculated in kg/tonne of fuel using specific fuel consumption.

BFO –Bunker Fuel Oil, MDO –Marine Diesel Oil, MGO –Marine Gas Oil

BC fraction of PM (f-BC); BFO: 0.12, MDO/MGO: 0.31. Source: for further information see Appendix A

Table 4-13: Engine type weightings from the EMEP/EEA air pollutant emission inventory guidebook 2016 (appears as Table 3-7 in guidebook)

Ship category	SSD MDO /MGO	SSD BFO	MSD MDO /MGO	MSD BFO	HSD MDO /MGO	HSD BFO	GT MDO /MGO	GT BFO	ST MDO /MGO	ST BFO
Liquid bulk ships	0.87	74.08	3.17	20.47	0.52	0.75	0.00	0.14	0.00	0.00
Dry bulk carriers	0.37	91.63	0.63	7.29	0.06	0.02	0.00	0.00	0.00	0.00
Container	1.23	92.98	0.11	5.56	0.03	0.09	0.00	0.00	0.00	0.00
General cargo	0.36	44.59	8.48	41.71	4.30	0.45	0.00	0.10	0.00	0.00
Ro Ro Cargo	0.17	20.09	9.86	59.82	5.57	2.23	2.27	0.00	0.00	0.00
Passenger	0.00	3.81	5.68	76.98	3.68	1.76	4.79	3.29	0.00	0.02
Fishing	0.00	0.00	84.42	3.82	11.76	0.00	0.00	0.00	0.00	0.00
Others	0.48	30.14	29.54	19.63	16.67	2.96	0.38	0.20	0.00	0.00
Tugs	0.00	0.00	39.99	6.14	52.80	0.78	0.28	0.00	0.00	0.00

SSD - Slow Speed Diesel, MSD – Medium Speed Diesel, HSD - High Speed Diesel, GT – Gas Turbine,

ST – Steam Turbine; MDO –Marine Diesel Oil, MGO –Marine Gas Oil, BFO –Bunker Fuel Oil

Source: Trozzi, 2010

Table 4-14: Main engine power from EMEP/EEA air pollutant emission inventory guidebook 2016 (appears as Table 3-6 in guidebook)

Ship category	Main engine power (kW)	
	1997 fleet	2010 fleet
Liquid bulk ships	6.695	6.543
Dry bulk carriers	8.032	4.397
Container	22.929	14.871
General cargo	2.657	2.555
Ro Ro Cargo	7.898	4.194
Passenger	3.885	10.196
Fishing	837	734
Other	2.778	2.469
Tug	2.059	2.033

Table 4-15: Average speeds from the EMEP/EEA air pollutant emission inventory guidebook 2016 (appears as Table 3-14 in guidebook)

Ship Type	Ave. Cruise Speed (km/h)	Manoeuvring time (hours)	Hotelling time (hours)
Liquid bulk ships	26	1.0	38
Dry bulk carriers	26	1.0	52
Container	36	1.0	14
General Cargo	23	1.0	39
Ro-Ro Cargo	27	1.0	15
Passenger	39	0.8	14
Fishing	25	0.7	60
Other	20	1.0	27

Source: Elaboration from Entec (2002)

These calculations allow the generation of the following activity data shown in **Table 4-16** and **Table 4-17**.

Non-bunkering can be subtracted from the total to give the total for bunkering and off-port calls.

Table 4-16: Activity data for all traffic

All traffic	Total distance travelled ('000 km)		Number of journeys		Fuel-oil consumption (kt)		Gas-oil consumption (kt)	
	To Gibraltar	From Gibraltar	To Gibraltar	From Gibraltar	To Gibraltar	From Gibraltar	To Gibraltar	From Gibraltar
Liquid bulk ships	5,015	6,422	2,671	2,671	240.7	308.2	11.5	14.7
Dry bulk carriers	19,803	18,719	4,236	4,236	651.3	615.6	7.0	6.6
Container	188	353	144	144	15.0	28.3	0.2	0.4
General Cargo	2,771	3,467	1,045	1,045	54.9	68.7	8.2	10.2
Ro Ro Cargo	44	67	23	23	1.2	1.8	0.3	0.4
Passenger	224	262	315	315	11.0	12.9	1.9	2.3
Fishing	0	24	2	2	0.0	0.0	0.0	0.1
Other	1,538	1,605	763	763	20.5	21.4	18.2	19.0
Tug	83	374	352	352	0.1	0.6	1.6	7.2
Total	29,667	31,295	9,551	9,551	994.7	1,057.5	48.8	60.9

Table 4-17: Activity data for non-bunkering and in-port traffic only

Non-bunkering port traffic only	Total distance travelled (‘000 km)		Number of journeys		Fuel-oil consumption (kt)		Gas-oil consumption (kt)	
	To Gibraltar	From Gibraltar	To Gibraltar	From Gibraltar	To Gibraltar	From Gibraltar	To Gibraltar	From Gibraltar
Liquid bulk ships	345	945	953	953	16.6	45.4	0.8	2.2
Dry bulk carriers	210	277	124	124	6.9	9.1	0.1	0.1
Container	36	53	42	42	2.9	4.2	0.0	0.1
General Cargo	127	94	47	47	2.5	1.9	0.4	0.3
Ro Ro Cargo	2	1	4	4	0.0	0.0	0.0	0.0
Passenger	203	240	303	303	10.0	11.8	1.7	2.1
Fishing	-	-	-	-	-	-	-	-
Other	51	44	35	35	0.7	0.6	0.6	0.5
Tug	26	229	312	312	0.0	0.3	0.5	4.4
Total	1,000	1,884	1,820	1,820	39.6	73.3	4.1	9.6

4.2.3.4 Determining emissions

With the fuel use activity data, for both gas oil (marine diesel oil) and fuel oil (bunkers fuel oil), NAEI emission factors for the use of gas oil and fuel oil in shipping were applied to calculate emissions from the relevant pollutants, shown in **Table 4-18**.

Table 4-18: Emission factors used for the shipping inventory

Fuel type	Emission factors (kt/Mt fuel)		
	CO ₂	CH ₄	N ₂ O
Fuel oil	3,114	0.06	0.15
Gas oil	3,206	0.03	0.14


Only emissions from ship departures are included in the inventory as per GPC guidelines. Emissions from activities that are not attributable to Gibraltar (i.e. those that have been deemed 'out of scope' due to the purpose of their call, as shown in Table 4-11) are reported in 'Other Scope 3' and are therefore not included in BASIC or BASIC+ inventory totals.

4.2.4 Aviation

Gibraltar is served by its own airport located within its boundary (**Figure 4-10**). There is also a nearby airport at Malaga, which is sometimes used when flights are diverted or as an alternative to flying directly into Gibraltar. However, no information is available for Malaga airport so Gibraltar Airport only is included here. Emissions are estimated for the 'landing/take-off cycle' (LTO) and cruise phases of flights. Only departing aircraft are included in the inventory as per GPC guidance.


Aviation emissions from the cruise phase and LTO are allocated to scope 3 (transboundary).

Figure 4-10: Gibraltar's International Airport



4.2.4.1 Overview

Figure 4-11 gives a brief overview of how estimates of emissions due to aviation have been made; a more detailed explanation is given in the following sections.

Figure 4-11: Aviation methodology flow diagram

4.2.4.2 Raw data

The raw data for estimating emissions have come from the International Civil Aviation Organization (ICAO) via the Department for Transport (DfT). The data provide a detailed log of all the journeys between Gibraltar and UK and non-UK airports.

4.2.4.3 Determining activity

The aircraft that operated between Gibraltar and the UK in 2017 were the Airbus A320 and A319. A smaller turboprop aircraft, the ATR 72, operated on flights between Gibraltar and Tangiers in Morocco. The UK airports that operated flights to and from Gibraltar in 2017 were: Birmingham, Bristol, Gatwick, Heathrow, Luton and Manchester.

Flight distances are calculated from great circle distances between airport pairs uplifted by 9.5% to allow for aircraft flying non-direct routes, in accordance with IPCC guidance. Cruise emission factors (based on fuel consumption) are selected from the EMEP/EEA air pollutant emission inventory guidebook 2013 by interpolating between the standard flight distances presented.

Fuel consumption for an Airbus A319, Airbus A320 and an ATR 72 from the EMEP/EEA air pollutant emission inventory guidebook 2013 aviation annex¹⁶ are shown in **Table 4-19**.

Table 4-19: Illustrative dataset from the EMEP/EEA air pollutant emission inventory guidebook 2013

Fuel (kg)	Phase of flight	Standard flight distances (nm)						(1nm = 1.852 km)	
		125	250	500	750	1,000	1,500	2,000	
A319	Climb/cruise/descent	1074.6	1812.2	2819.8	3905.4	5013.8	7294.7	9513.4	
	LTO	688.8	688.8	688.8	688.8	688.8	688.8	688.8	
	Total	1763.4	2501.0	3508.7	4594.2	5702.6	7983.5	10202.2	

¹⁶ www.eea.europa.eu/publications/emep-eea-guidebook-2013/part-b-sectoral-guidance-chapters/1-energy/1-a-combustion/1-a-3-a-aviation-annex

A320	Climb/cruise/descent	1062.6	1831.1	2949.3	4200.4	5439.0	7994.8	10464.9
	LTO	873.3	873.3	873.3	873.3	873.3	873.3	873.3
	Total	1935.8	2704.4	3822.6	5073.6	6312.3	8868.1	11338.2
AT72	Climb/cruise/descent	723.5	1361.9	2652.5	3967.0	5276.0	8090.4	
	LTO	393.8	393.8	393.8	393.8	393.8	393.8	
	Total	1117.4	1755.7	3046.3	4360.9	5669.8	8484.2	

The 2016 EMEP/EEA air pollutant emission inventory guidebook provides a spreadsheet tool to calculate fuel consumptions and emissions during the LTO cycle. This tool includes airport specific taxiing times by year. The latest year available for Gibraltar is 2015, which gave taxi-out and taxi-in times of 605 s and 204 s, respectively. These times, along with the aircraft fleet mix, have been used to calculate LTO emissions for 2017.

4.2.4.4 Determining emissions

The calculation for emissions is shown in **Equation 4-2**.

Equation 4-2: Equation for aviation emission estimation

$$Emissions = \sum (LTO \text{ fuel use} * LTO \text{ EF}), (Cruise \text{ fuel use} * fuel \text{ EF})$$

LTO = Landing/take-off cycle

EF = Emission factor

To generate total fuel consumption, the total number of flights broken down by destination airport and aircraft type were multiplied by the emission factors, interpolated on distance, from **Table 4-19**. These were then summed to give the values **Table 4-20**.

The emission factors in **Table 4-21** were then used to calculate total emissions. The fuel use factors assume jet kerosene from Tables 3.6.4 and 3.6.5 of the 2006 IPCC Guidelines; Emission factors for methane for LTO cycle are taken from Table 3-5 in the EMEP/EEA air pollutant emission inventory guidebook 2013 for an Airbus A320: examples of aircraft types and emission factors for LTO cycles as well as fuel consumption per aircraft type, kg/LTO.

It is assumed that emissions from all aircraft departing Gibraltar Airport are allocated to Gibraltar. This is because, although there may be some use of the airport by non-Gibraltarian residents/visitors, these numbers are impossible to determine with any accuracy, it is assumed the majority of visitors arriving at Gibraltar Airport are likely to be resident or visiting.

LTO cycle emissions include emissions from both take-off at the departure airport and landing at the destination airport. However, for each departure from Gibraltar there is an associated arrival movement at Gibraltar that has emissions that are equivalent to the emissions from landing at the destination airport. Therefore, counting all the LTO cycle emissions associated with departures from Gibraltar captures all the LTO cycle emissions at the airport. For cruise only the departure emissions are included in the inventory for Gibraltar.

Table 4-20: Total annual fuel consumption by aircraft

Phase of flight	Unit	Domestic (UK)	International (non-UK)
LTOs	Number	2,111	105
Total fuel consumption	Kt	12.79	0.04
Of which cruise	Kt	11.45	0.02
Total fuel consumption	TJ	561.50	1.69
Of which cruise	TJ	502.56	1.02

Table 4-21: Emission factors for aircraft phases by pollutant

Phase of flight	Pollutant	Unit	Emission factor
Cruise	Carbon	kt/TJ fuel	0.0195
Cruise	CH ₄	kt/TJ fuel	Zero
Cruise	N ₂ O	kt/TJ fuel	0.000002
LTO	Carbon	kt/LTO	0.00070909
LTO	CH ₄	kt/LTO	0.0000001
LTO	N ₂ O	kt/LTO	0.0000001

4.3 Waste

The waste profile of Gibraltar is unique due to the territory's location, restricted land area, high population density and absence of heavy industry. The majority of waste generated in Gibraltar is municipal, largely arising at households and commercial premises.

This report has applied the methodologies recommended under the GPC Guidelines for the estimation of GHG emissions from waste. Where possible, quantities of CO₂, CH₄ and N₂O have been estimated from the following sources based on activities during 2017:

- Solid waste disposal.
- Biological treatment of solid waste.
- Incineration.
- Wastewater.

Emissions from waste are allocated by scope to the location they are emitted.


Therefore, the treatment of biological waste, MSW and some incineration in Spain are allocated to scope 3.

Emissions from incineration in Gibraltar are scope 1.

Emissions from wastewater are out of boundary so allocated to scope 3.

4.3.1 Summary

Figure 4-12 gives a brief overview of how waste emissions have been estimated, with a more detailed explanation provided in the following sections.

Figure 4-12: Waste methodology flow diagram

4.3.2 Raw data

4.3.2.1 Municipal solid waste

Municipal waste, generated at households, commercial premises and state-run facilities, such as schools and hospitals, is collected six days a week by a waste management contractor. This waste is then transported in bulk to the Complejo Medioambiental, Sur de Europa, in Los Barrios, Spain via a temporary waste transfer station in Gibraltar. At Los Barrios, waste is manually and mechanically sorted to remove the recyclable fraction. Biological waste is also removed for composting and the remaining fraction is disposed of to landfill.

Data on the total quantity of MSW arisings by weight for Gibraltar have been provided by the Government of Gibraltar, as shown in **Table 4-22**.

Table 4-22: MSW arisings in Gibraltar in 2017

Month	Household (tonnes)	Refuse (tonnes)	Total (tonnes)
January 2017	721	1,129	1,850
February 2017	946	1,277	2,223
March 2017	1,018	1,623	2,641
April 2017	527	1,261	1,788
May 2017	778	1,685	2,463
June 2017	1,235	1,399	2,634
July 2017	966	1,400	2,367
August 2017	1,064	1,623	2,687
September 2017	1,100	876	1,976
October 2017	1,141	1,644	2,785
November 2017	1,183	1,485	2,668
December 2017	889	1,297	2,186
Total	11,568.76	16,698.22	28,267

4.3.2.2 Clinical waste

Gibraltar's clinical waste is generated by a number of sources including dental and veterinary practices, and medical premises. In 2008, a new incinerator was commissioned in Gibraltar for the sole purpose of treating clinical waste. Although the incinerator has adequate capacity for the treatment of all clinical waste arisings within the boundary, maintenance issues will occasionally result in clinical waste being exported to an incinerator in Spain.

Details on the quantity of clinical waste incinerated within Gibraltar in 2017 are provided by the Gibraltar Health Authority. Volumes of clinical waste incinerated are based on average bin weight of 7.5Kg per 60 litre bin of waste. The Government of Gibraltar provides information regarding the amount of clinical waste exported to Spain for incineration.

4.3.3 Determining activity

4.3.3.1 Composition of MSW

To determine the fraction of degradable organic carbon (DOC), the composition of MSW arisings have been estimated by applying the results of the 2014 Waste Characterisation Study to the total reported MSW detailed above. The study was completed by the Department of the Environment. It analysed MSW from three collection routes, in March and August 2014, in Gibraltar recording the waste type, weight and bulk density.

The waste categories have been grouped into three assumed treated groups; biological treatment (composted), landfill and recycled. A summary of the results and the treatment groups are provided in **Table 4-23**.

Table 4-23: Results of the 2014 Waste Characterisation Study and assumed treatment groups

<i>Waste category</i>	Average waste composition	Assumed treatment route
	Weight (%)	
Paper & Cardboard	25.1	Recycled
Dense Plastics	7.0	Recycled
Plastic Film	6.1	Landfill
Organics	30.7	Composted
Metals	3.4	Recycled
Glass	4.9	Recycled
Composites	2.2	Landfill
Special Municipal waste	3.0	Landfill
Textiles	3.2	Recycled
Fines	0.5	Landfill
Unclassified Combustibles	12.4	Landfill
Unclassified Incombustibles	0.9	Landfill
WEEE	0.5	Recycled
Batteries	0.0	Recycled
Total	100	

4.3.4 Determining emissions

4.3.4.1 Solid waste disposal

Emissions of CH₄ from landfilling MSW have been calculated using the ‘Methane Commitment’ method. This allows emissions to be estimated based on the quantity of waste sent to landfill in a single year by adopting a mass balance approach. Prior to this, it was necessary to determine the ‘methane generation potential’ of the waste landfilled. The formulas for each are provided in formulas 8.1, 8.3 and 8.4 of the GPC, as below (**Figure 4-13**). A change was made to the DOC value to exclude waste categories that we assume are not landfilled (e.g. paper/card, food, etc.), and include waste categories that are landfilled (e.g. nappies). The DOC value used in the calculation is therefore 0.246 tonnes C/tonne waste.

Figure 4-13: GPC equations for calculating emissions from landfill

Equation 8.3 Methane commitment estimate for solid waste sent to landfill

$$CH_4 \text{ emissions} = MSW_x \times L_0 \times (1-f_{rec}) \times (1-OX)$$

Description		Value
CH ₄ emissions	= Total CH ₄ emissions in metric tonnes	Computed
MSW _x	= Mass of solid waste sent to landfill in inventory year, measured in metric tonnes	User input
L ₀	= Methane generation potential	Equation 8.4 Methane generation potential
f _{rec}	= Fraction of methane recovered at the landfill (flared or energy recovery)	User input
OX	= Oxidation factor	0.1 for well-managed landfills; 0 for unmanaged landfills

Source: Adapted from Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories

Equation 8.4 Methane generation potential, L₀

$$L_0 = MCF \times DOC \times DOC_f \times F \times 16/12$$

Description		Value
L ₀	= Methane generation potential	Computed
MCF	= Methane correction factor based on type of landfill site for the year of deposition (managed, unmanaged, etc., fraction)	Managed = 1.0 Unmanaged (≥5 m deep) = 0.8 Unmanaged (<5 m deep) = 0.4 Uncategorized = 0.6
DOC	= Degradable organic carbon in year of deposition, fraction (tonnes C/tonnes waste)	Equation 8.1
DOC _f	= Fraction of DOC that is ultimately degraded (reflects the fact that some organic carbon does not degrade)	Assumed equal to 0.6
F	= Fraction of methane in landfill gas	Default range 0.4-0.6 (usually taken to be 0.5)
16/12	= Stoichiometric ratio between methane and carbon	

Source: IPCC Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories (2000)

Equation 8.1 Degradable organic carbon (DOC)⁵²

$$\text{DOC} = (0.15 \times A) + (0.2 \times B) + (0.4 \times C) + (0.43 \times D) + (0.24 \times E) + (0.15 \times F)$$

A	=	Fraction of solid waste that is food
B	=	Fraction of solid waste that is garden waste and other plant debris
C	=	Fraction of solid waste that is paper
D	=	Fraction of solid waste that is wood
E	=	Fraction of solid waste that is textiles
F	=	Fraction of solid waste that is industrial waste

Source: GPC

4.3.4.2 Biological treatment

The emissions of CH₄ and N₂O from the biological treatment of waste have been calculated using equation 8.5 from the GPC guidelines (**Figure 4-14**) and emission factors for composting given in the GPC; these are detailed in **Table 4-24**. It is assumed that waste is treated whilst wet, as we have no information on whether waste is dried before being treated. This year a revised emission factor for N₂O from composting has been applied from the IPCC 2006 Guidelines Waste Chapter update.

As the Los Barrios waste treatment facility only provides composting as a form of biological treatment, it has been assumed this is the sole method of biological treatment.

Table 4-24: Biological waste treatment emission factors

GHG	Emission factor
CH ₄	4g per kg of wet waste treated
N ₂ O	0.24g per kg of wet waste treated

Figure 4-14: GPC equation for calculating emissions from biological treatment of waste

Equation 8.5 Direct emissions from biologically treated solid waste

$$\text{CH}_4 \text{ Emissions} = (\sum_i (m_i \times F_{\text{CH}_4}) \times 10^{-3} - R)$$

$$\text{N}_2\text{O Emissions} = (\sum_i (m_i \times \text{EF}_{\text{N}_2\text{O}_i}) \times 10^{-3})$$

Description	Value
CH ₄ emissions	– Total CH ₄ emissions in tonnes Computed
N ₂ O emissions	– Total N ₂ O emissions in tonnes Computed
m	– Mass of organic waste treated by biological treatment type i, kg User input
EF _{CH4}	– CH ₄ emissions factor based upon treatment type, i User input or default value from table 8.3 Biological treatment emission factor
EF _{N2O}	– N ₂ O emissions factor based upon treatment type, i User input or default value from table 8.3 Biological treatment emission factor
i	– Treatment type: composting or anaerobic digestion User input
R	– Total tonnes of CH ₄ recovered in the inventory year, if gas recovery system is in place User input, measured at recovery point

Source: 2006 IPCC Guidelines for National Greenhouse Gas Inventories, Volume 5, Chapter 4: Biological Treatment of Solid Waste

Source: GPC

4.3.4.3 Clinical waste incineration

The emission of CH₄ and N₂O from the incineration of clinical waste has been calculated using emission factors provided in the UK NAEI 2016. The emission factors are provided in **Table 4-25**.

Table 4-25: Clinical waste incineration emission factors

GHG	Emission factor	Unit
CH ₄	240	kt/mt waste incinerated
N ₂ O	0.02	kt/mt waste incinerated
CO ₂	0.03	kt/mt waste incinerated

4.3.5 Wastewater

Wastewater in Gibraltar is pumped out to sea with no treatment. However, the Gibraltar Government has awarded an Advanced Works Contract to the joint venture between NWG Commercial Services Limited [Northumbrian Water] and Modern Water to design, construct, operate and maintain a wastewater treatment facility in Gibraltar. This is planned to be operational by 2020.

Emissions from pumping are reported under stationary combustion scope 2 emissions (consumption of electricity). Emissions from wastewater have been calculated by scaling UK data. These are:

- Biochemical oxygen demand (BOD) and nitrogen content on a per person per day basis.
- Tonnes of N₂O per million people.

The IPCC CH₄ conversion factor for wastewater to sea/lakes/rivers was used to estimate CH₄ – this is also given in the GPC. This is likely to overestimate emissions as it assumes anaerobic decomposition in stagnant water, and ocean decomposition is likely to be much less stagnant and, therefore, undergoes higher aerobic decomposition with lower associated emissions. N₂O emission assumptions do not account for denitrification in sewage treatment or alternative disposal methods (e.g. to land, incineration). It is assumed that all sewage is discharged in raw form to sea.

The equations for calculating emissions from wastewater are given below.

Figure 4-15: GPC equations for calculating emissions from wastewater treatment

Equation 8.9 CH₄ generation from wastewater treatment

$\text{CH}_4 \text{ emissions} = \sum_i [(TOW_i - S_i) EF_i - R_i] \times 10^{-3}$		
Description		Value
CH ₄ emissions	= Total CH ₄ emissions in metric tonnes	Computed
TOW _i	= Organic content in the wastewater For domestic wastewater: total organics in wastewater in inventory year, kg BOD/yr ^{Note 1} For industrial wastewater: total organically degradable material in wastewater from industry i in inventory year, kg COD/yr	Equation 8.10
EF _i	= Emission factor kg CH ₄ per kg BOD or kg CH ₄ per kg COD ^{Note 2}	Equation 8.10
S _i	= Organic component removed as sludge in inventory year, kg COD/yr or kg BOD/yr	User input
R _i	= Amount of CH ₄ recovered in inventory year, kg CH ₄ /yr	User input
i	= Type of wastewater For domestic wastewater: income group for each wastewater treatment and handling system For industrial wastewater: total organically degradable material in wastewater from industry i in inventory year, kg COD/yr	Equation 8.10

Note 1: Biochemical Oxygen Demand (BOD): The BOD concentration indicates only the amount of carbon that is aerobically biodegradable. The standard measurement for BOD is a 5-day test, denoted as BOD₅. The term "BOD" in this chapter refers to BOD₅.

Note 2: Chemical Oxygen Demand (COD): COD measures the total material available for chemical oxidation (both biodegradable and non-biodegradable).

Source: 2006 IPCC Guidelines for National Greenhouse Gas Inventories, Volume 5, chapter 6: Wastewater Treatment and Discharge

Equation 8.10 Organic content and emission factors in domestic wastewater⁵⁵

$$TOW_i = P \times BOD \times I \times 365$$

$$EF_j = B_o \times MCF_j \times U_i \times T_{i,j}$$

Description	Value
TOW_i = For domestic wastewater: total organics in wastewater in inventory year, kg BOD/yr	Computed
P = City's population in inventory year (person)	User input ⁵⁶
BOD = City-specific per capita BOD in inventory year, g/person/day	User input
I = Correction factor for additional industrial BOD discharged into sewers	In the absence of expert judgment, a city may apply default value 1.25 for collected wastewater, and 1.00 for uncollected. ⁵⁷
EF_j = Emission factor for each treatment and handling system	Computed
B_o = Maximum CH ₄ producing capacity	User input or default value: • 0.6 kg CH ₄ /kg BOD • 0.25 kg CH ₄ /kg COD
MCF_j = Methane correction factor (fraction)	User input ⁵⁸
U_i = Fraction of population in income group i in inventory year	
$T_{i,j}$ = Degree of utilization (ratio) of treatment/discharge pathway or system, j, for each income group fraction i in inventory year	User input ⁵⁹

Source: 2006 IPCC Guidelines for National Greenhouse Gas Inventories, Volume 5, chapter 6: Wastewater Treatment and Discharge

Equation 8.11 Indirect N₂O emissions from wastewater effluent

$$N_2O \text{ emissions} = [(P \times \text{Protein} \times F_{NPR} \times F_{NON-CON} \times F_{IND-COM}) - N_{SLUDGE}] \times EF_{EFFLUENT} \times 44/28 \times 10^{-3}$$

Description	Value
N ₂ O emissions = Total N ₂ O emissions in tonnes	Computed
P = Total population served by the water treatment plant	User input
Protein = Annual per capita protein consumption, kg/person/yr	User input
$F_{NON-CON}$ = Factor to adjust for non-consumed protein	1.1 for countries with no garbage disposals, 1.4 for countries with garbage disposals
F_{NPR} = Fraction of nitrogen in protein	0.16, kg N/kg protein
$F_{IND-COM}$ = Factor for industrial and commercial co-discharged protein into the sewer system	1.25
N_{SLUDGE} = Nitrogen removed with sludge, kg N/yr	User input or default value: 0
$EF_{EFFLUENT}$ = Emission factor for N ₂ O emissions from discharged to wastewater in kg N ₂ O-N per kg N ₂ O	0.005
44/ 28 = The conversion of kg N ₂ O-N into kg N ₂ O	

Source: 2006 IPCC Guidelines for National Greenhouse Gas Inventories, Volume 5, chapter 6: Wastewater Treatment and Discharge

4.4 Industrial Processes and Product Use (IPPU)

The industrial processes and product use (IPPU) sector covers GHG emissions from a range of activities. The main emission sources are releases from industrial processes that chemically or physically transform materials (e.g. blast furnaces in the iron and steel industry, and ammonia and other chemical products manufactured from fossil fuels used as chemical feedstock). During these processes, many different GHGs, including CO₂, CH₄, N₂O, hydrofluorocarbons (HFCs) and perfluorocarbons (PFCs) can be produced. Emissions also occur from the use of products such as solvents, aerosols and inhalers, and anaesthetics.

4.4.1 Summary

Industrial processes specifically covered by the GPC include:

- Production and use of mineral products.
- Production and use of chemicals.
- Production of metals.

None of these activities occur in Gibraltar, so this source is NO.

Product use in the GPC covers:

- Lubricants and paraffin waxes used in non-energy products.
- HFC gases used in electronics production.
- Fluorinated gases used as substitutes for ozone-depleting substances.

4.4.2 Separating IPPU GHG emissions and energy-related GHG emissions

Allocation of emissions from the use of fossil fuels between the stationary energy and IPPU sectors can be complex. The GPC follows IPCC Guidelines¹⁷, which define 'fuel combustion' in an industrial process context as 'the intentional oxidation of material within an apparatus that is designed to provide heat or mechanical work to a process, or for use away from the apparatus.'

Therefore:

- If the fuels are combusted for energy use, the emission from fuel uses shall be counted under stationary energy.
- If the derived fuels are transferred for combustion in another source category, the emissions shall be reported under stationary energy.
- If combustion emissions from fuels are obtained directly or indirectly from the feedstock, those emissions shall be allocated to IPPU.
- If heat is released from a chemical reaction, the emissions from that chemical reaction shall be reported as an industrial process in IPPU.

In the case of Gibraltar, in the stationary combustion category all fuels are combusted for energy use so emissions are accounted for in this sector and not IPPU.

4.4.3 Determining activity

The industrial processes identified above are NO, so no data are available.

N₂O emissions from medical anaesthetics have been estimated using delivery information supplied by the hospital's medical gas supplier.

HFC emissions from metered dose inhalers (MDIs) have been estimated using information supplied by the Gibraltar Health Authority regarding the total number of MDIs prescribed in Gibraltar in 2017.

In product use, emissions of fluorinated gases (the so-called F-gases) have been estimated based on a scaling of UK data using an appropriate indicator. The source categories of these emissions and the indicators used are shown in **Table 4-26**.

¹⁷ Box 1-1 from 2006 IPCC Guidelines for National Greenhouse Gas Inventories, Volume 3 IPPU, Chapter 1 introduction.

Table 4-26: F-gas emission sources and activities

Source	Activity	Indicator
Aerosols - halocarbons	non-fuel combustion	Population
Firefighting	non-fuel combustion	GDP
Foams	non-fuel combustion	GDP
Commercial refrigeration	Refrigeration and air-conditioning - lifetime	GDP
Domestic refrigeration	Refrigeration and air-conditioning - lifetime	Population
Industrial refrigeration	Refrigeration and air-conditioning - lifetime	GDP
Mobile Air Conditioning	Refrigeration and air-conditioning - lifetime	Number of vehicles
Refrigerated Transport	Refrigeration and air-conditioning - lifetime	GDP
Stationary Air Conditioning	Refrigeration and air-conditioning - lifetime	Population
Electrical Insulation	non-fuel combustion	GDP
Precision cleaning	non-fuel combustion	GDP

*Gross domestic product (GDP)

4.4.4 Estimating emissions

Emissions have been estimated by multiplying the factor for the UK by the associated indicator for Gibraltar (GDP, population, etc.).

Estimates of N₂O emissions from anaesthetics have been calculated using an emission factor of 1 as it is assumed that none of the administered N₂O is chemically changed by the body, and all is returned to the atmosphere, so therefore, it is reasonable to assume an emission factor of 1.0 (IPCC 2006 GL).

The emissions factor used is based on an assumption that each MDI contains 12g of HFC per MDI¹⁸. The split of HFCs was calculated using UK NAEI assumption that 96% of MDIs are formulated with HFC-134a and 4% are formulated with HFC-227ea.

Emissions from the use of electrical insulation and precision cleaning have been added for the 2017 inventory. Recalculations have also been made to add these emissions to the 2015 and 2016 inventories.

4.5 Agriculture, Forestry, and Other Land Use (AFOLU)

Gibraltar has no notable agriculture, so this emission source has not been estimated, and is noted as 'NO'.

Gibraltar is also regarded as having no emissions from land use, land use change and forestry (LULUCF), so this emission source is also noted as 'NO'.

¹⁸ Gluckman (2013). NAEI – Report on F-Gases. Report on Programme of Work on F-Gases, Financial Year 2013/14. Version 2, November 11th 2013. Report prepared by Ray Gluckman, SKM Enviro, Sinclair Knight Merz, New City Court, 20 St Thomas Street, London, SE1 9R, UK

5 Results

This section sets out the results of the Gibraltar city GHG inventory. As detailed in the methodology section above, this inventory considered all sources attributable to Gibraltar, following the methods published by the GPC guidelines.

The inclusion of different sources in the reported total varies by chosen reporting standard, and these sources also vary in their level of potential influence: Gibraltar has limited control over the emissions associated with water-borne navigation, for example, whereas power generation can be much more easily affected through local decision-making. Water-borne navigation emissions dominate the results, discussed below, and overshadow other sources for which Gibraltar has more influence. With this in mind, the results section presents the results both as a whole – total emissions for Gibraltar across all sources – and distinguishes between different reporting levels and sources, including presenting a subset of BASIC+ emissions (the recommended reporting level) termed the ‘manageable emissions’, excluding international transboundary shipping.

5.1 Summary

Total emissions for Gibraltar in 2017 by different reporting level are shown in **Figure 5-1**. Sources included within each reporting level are detailed in **Table 5-1** and summarised in **Table 5-2** below. Emissions included within each higher reporting level are cumulative from lower levels. It is current best practice for cities to report BASIC+ emissions wherever possible, and this chart represents emission sources as classified by the GPC.

Figure 5-1 – Emissions disaggregated by scope and by reporting level (BASIC and BASIC+)




Table 5-1 - Classification of emission categories by scope and reporting level. Note, these are cumulative, and higher reporting levels include those sources in lower levels

Scope	BASIC	BASIC+	Outside of scopes
Scope 1	Emissions from in-boundary fuel combustion Emissions from in-boundary production of energy used in auxiliary operations In-boundary fugitive emissions	In-boundary emissions from industrial processes In-boundary emissions from product use In-boundary emissions from livestock In-boundary emissions from land	

Scope	BASIC	BASIC+	Outside of scopes
	Emissions from in-boundary transport Emissions from waste and wastewater generated and treated within the city	In-boundary emissions from other agriculture	
Scope 2	Emissions from consumption of grid-supplied energy		
Scope 3	Emissions from waste and wastewater generated within but treated outside of the city	Transmission and distribution losses from grid-supplied energy Emissions from transboundary journeys	
Outside of scopes			Electricity generation ¹ International bunkers Vehicle fuel exports

¹ Reported for information only. Electricity emissions are allocated to the end-user.

Figure 5-2 – Gibraltar’s ‘manageable’ emissions by source category for 2017 (under the GPC’s BASIC+ reporting, excluding transboundary waterborne navigation and other scope 3)




Figure 5-3 – Gibraltar 's total emissions (including excluded sources) by source category for 2017.




Table 5-2 – Total emissions for Gibraltar in 2017 by source.

Sector	Sub-sector	Total GHGs (metric tonnes CO ₂ e)			
		Scope 1	Scope 2	Scope 3	Total
Stationary Energy	Residential buildings	NO	43,348	9,085	52,433
	Commercial and institutional buildings and facilities	1,423	81,477	17,075	99,976
	Manufacturing industries and construction	NO	NO	NO	
	Energy industries	NO	2,681	IE	2,681
	Energy generation supplied to the grid	154,317			
	Agriculture, forestry and fishing activities	NO	NO	NO	
	Non-specified sources	NO	538	113	651
	Fugitive emissions from mining, processing, storage, and transportation of coal	NO			
	Fugitive emissions from oil and natural gas systems	NO			
	SUBTOTAL		1,423	128,044	26,273
Transport	On-road transportation	55,438	NO	IE	55,438
	Railways	NO	NO	NO	
	Waterborne navigation	12,576	NO	262,783	275,359
	Aviation	NO	NO	38,482	38,482
	Off-road transportation	IE	NO	IE	
	SUBTOTAL		68,014		301,265
Waste	Solid waste generated in the city	NO		15,782	15,782

Sector	Sub-sector	Total GHGs (metric tonnes CO ₂ e)			
		Scope 1	Scope 2	Scope 3	Total
	Biological waste generated in the city	NO		1,489	1,489
	Incinerated and burned waste generated in the city	345		54	398
	Wastewater generated in the city	NO		1,790	1,790
	Solid waste generated outside the city	NO			
	Biological waste generated outside the city	NO			
	Incinerated and burned waste generated outside city	NO			
	Wastewater generated outside the city	NO			
	SUBTOTAL	345		19,115	19,460
Industrial Processes and Product Use	Emissions from industrial processes occurring in the city boundary	NO			
	Emissions from product use occurring within the city boundary	10,591			10,591
	SUBTOTAL	10,591			10,591
Other Scope 3	SUBTOTAL			3,324,642	3,324,642
TOTAL		80,372	128,044	3,671,294	3,879,711

Note: Agriculture, Forestry, and Other Land Use emissions are not estimated within this inventory and are considered negligible

Colour coding of Table 5.2

	BASIC sources
	BASIC+ sources
	Additional scope 1 sources required for territorial reporting
	Other scope 3 sources

5.2 Total emissions for Gibraltar

Total emissions for Gibraltar, from all calculated sources are presented in **Table 5-2** and **Figure 5-3** above.

Overall, the largest contributor of emissions to the Gibraltar city inventory is 'Other scope 3' accounting for 86% of emissions. 'Other scope 3' is dominated by marine bunkering (99%), with a small contribution (1%) from non-Gibraltarian road transport emissions. Transboundary transportation sources are included in BASIC+ reporting, but have been excluded from the chart in **Figure 5-2** to better represent emissions attributable to and influenced by the community. Note that emissions from private boats are captured under Scope 1.

Stationary energy is responsible for 4% of total emissions, waste 0.5%, and industrial processes and product use (IPPU), 0.3%. Transport emissions from in-scope sources comprise 2.7% of total emissions, of which 75% are attributable to waterborne transport.

As **Table 5-2** illustrates, Scope 1 emissions are largely dominated by road transport fuel use, but there is also a noticeable contribution from hydrofluorocarbons (HFCs) from product use (such as air conditioning units).

Scope 2 emissions from electricity consumption are also large, due to the reliance on electricity for energy requirements and generation technologies. Because gas oil is used to generate electricity, the emissions per kilowatt hour (kWh) are considerably higher than, for example, those in the UK. The implied emission factor based on fuel consumption in power stations and total output in Gibraltar is 0.69kg CO₂e/kWh, compared to the UK grid factor of 0.35kg CO₂e/kWh¹⁹. The implied emission factor for Gibraltar in the 2016 inventory was 0.91 kg CO₂e/kWh and 0.86 kg CO₂e/kWh for 2015.

Scope 3 emissions are largest overall across scopes, due primarily to shipping activities and bunkering.

Table 5-3 – Emissions by sector and reporting standard

Sector	BASIC	BASIC+	Manageable	BASIC+ and Scope 3
Stationary Energy	129,467	155,740	155,740	155,740
Transportation	68,014	369,279	106,496	369,279
Waste	19,460	19,460	19,460	19,460
IPPU		10,591	10,591	10,591
Other Scope 3				3,324,642
TOTAL	216,941	555,069	292,286	3,879,711

As **Table 5-3** illustrates, there is a large difference between the reported emissions between the GPC's BASIC and BASIC+ reporting levels. This is due to the inclusion of additional sources within BASIC+ which are significant within Gibraltar, namely transboundary (scope 3) emissions from shipping, and lesser contributions from aviation. Further inclusion of emissions that are deemed 'outside of scopes' (i.e. they are reported for information but are not deemed to be within the influence of responsibility of the city, such as bunkering fuels) dwarf the BASIC and BASIC+ when considered, contributing to 84% of total emissions.

5.3 Total manageable emissions

The recommended reporting approach for city-level emissions under the GPC is BASIC+, therefore excluding emissions from combustion of bunkering fuels. Transboundary transport emissions are included under BASIC+ reporting however, and this includes a large proportion of waterborne navigation emissions. This presents a particularly large source for Gibraltar, and is one that the community has little influence over. It also dominates the results, making it difficult to identify the impact

¹⁹ <https://www.gov.uk/government/publications/greenhouse-gas-reporting-conversion-factors-2017>


of smaller, more manageable local sources. For this reason, waterborne navigation (scope 3, transboundary) has been excluded from the total presented in **Figure 5-2**. Private marine emissions are retained. This subset, therefore, may be considered Gibraltar's 'manageable emissions'.

When these sources are removed, the inventory results are much more aligned to those expected for a city (see **Figure 5-4**), with stationary energy dominating, accounting for 53% of emissions. Transportation contributes 36%; 19% is attributable to road transport, 4% to waterborne navigation, and 13% to aviation. Contributions from waste and IPPU sectors are smaller, contributing 7% and 3% respectively.

Gibraltar's per capita emissions are 6.3 tCO_{2e}, based on the 'BASIC' emissions profile. This indicates that Gibraltar has slightly higher emissions per capita, compared with other cities and the UK average of 5.5 tonnes per person. In 2017 However, it is important to acknowledge Gibraltar's small resident population, its unique geographical situation compared to most global cities, and the impacts and limitations this places upon emissions. Cities with similar per capita emissions to Gibraltar include Boston and New Orleans²⁰.

Figure 5-4 - Sectoral breakdown of latest community GHG emission inventories from Carbonn Cities Climate Registry (cCCR) report²¹

Performance



5.4 Comparison with past inventories

This section aims to compare the 2017 inventory results against the revised 2016 (2016r) and 2015 (2015r) inventory inventories. There are some differences between the original 2015 inventory²², 2016 inventory²³ and the revised versions used as the comparison in this section; this is due to improvements in methodologies and activity data availability during the compilation of the 2017 inventory, which have been applied retrospectively to previous year's inventories for consistency and accuracy, following international best practice. Important recalculations are explained in **Appendix 3**. The 2017 inventory has not been compared to the 2013 inventory; the 2013 inventory was a 'pilot' using a pilot version of the GPC. For Gibraltar's city inventory programme, the 2015 inventory is the first official inventory. The 2013 inventory is also not directly comparable to the 2015-2017 inventories due to a large number of method changes and an updated reporting approach using the now-finalised GPC.

With only three directly comparable inventories, it is difficult to confidently comment on changes to the time series between the two years; however, this section highlights key changes and aims to provide

²⁰ https://www.c40.org/research/open_data/5

²¹ http://carbonn.org/fileadmin/user_upload/cCCR/ccr-digest-2014-2015/ccr-digest-2014-2015-online-final.pdf

²² https://www.gibraltar.gov.gi/new/sites/default/files/HMGoG_Documents/20170601-Gibraltar_City_Inventory_Report_Published.pdf

²³ https://www.gibraltar.gov.gi/new/sites/default/files/HMGoG_Documents/2016-GibraltarCityInventory_Report_Final.pdf

some insight and interpretation. As Gibraltar’s inventory programme progresses, there will be a longer time series of inventories, allowing more confident commentary on annual emission trends.


Emissions from the 2015r, 2016r and 2017 inventories are presented, by sector, in **Table 5-4** and **Figure 5-5**.

More information on the specific reasons for changes between the 2015r and 2016 inventories and a comparison of years by sector and sub-sector is found in **Appendix 2**. Information on the revisions between the 2015 and 2015r inventories is given in **Appendix 3**.

Table 5-4 – Comparison between the 2015r and 2016 inventories by scope

Reporting sector	Emissions (tCO ₂ e)		
	2015r	2016r	2017
Stationary Energy	193,439	183,707	155,740
<i>Transportation (aIP²⁴)</i>	283,488	376,318	369,279
Transportation (excluding scope 3 shipping)	93,589	103,035	106,496
Waste	22,249	21,561	19,460
IPPU	11,061	10,874	10,591
<i>Other Scope 3²⁴</i>	3,077,423	3,206,983	3,324,642
Total Manageable emissions	320,338	319,177	292,286

Figure 5-5 - Gibraltar's 'manageable' emissions for 2015r, 2016r and 2017



* Transport emissions excluding scope 3 shipping

²⁴ not included in Gibraltar's 'manageable' emissions

5.4.1 Summary of changes

Highlights

Gibraltar's total manageable emissions have decreased by 9% since 2015r and 8% since 2016r; this is a result of the following:

- ↓ Emissions from electricity generation have decreased by 19% since 2015r and 15% since 2016r; this is due to less fuel being used to generate a unit of electricity, implying improvements in efficiency at Gibraltar's electricity power stations. Electricity consumption by residents and activity in Gibraltar remained fairly consistent between 2015 and 2017.
- ↓ Emissions from Waste are around 13% and 10% lower in 2017 than 2015r and 2016r (respectively) due to a reduction in total waste arisings sent to landfill (and composting).
- ↓ Emissions from IPPU have decreased by 4% between 2015r and 2017; this follows trends in UK data that is used as a proxy for Gibraltar's emissions from product use (e.g. air conditioning and refrigeration).
- ↑ Emissions from road transport in Gibraltar have increased by 5% due to more fuel being consumed by vehicles in Gibraltar.
- ↑ Emissions from scope 3 waterborne navigation²⁵ are 44% higher in the 2016r inventory than the 2015r inventory, largely due to longer liquid bulk container departing journey lengths in 2016; the reason for this is currently unknown. Trends in other vessel categories were more stable between the two years. These emissions have declined slightly since 2016r.
- ↑ Emissions from aviation are around 36% higher in 2017 than 2015r, likely due to an increased number of flights to London Gatwick, London Heathrow and Manchester between 2015r and 2016r. Between 2016r and 2017, aviation emissions have increased by 4%, largely due to an increased number of flights to London Gatwick and Manchester.

Stationary Energy – Electricity


Although not presented in the scope 1 totals (to avoid double counting), emissions from the generation of electricity have decreased by 15% since 2016r and 20% since 2015r; this is due to considerably less fuel use for electricity generation at Gibraltar's power stations.

Scope 2 emissions in Gibraltar are solely those from **electricity consumption** and between 2015r and 2017, total scope 2 emissions have shown a decrease of over 30kt CO₂e or 19%. This decrease in emissions is despite total electricity consumption being fairly stable from 2015r to 2017, and is due to considerably less fuel being used at the power stations to generate a unit of electricity (**Figure 5-6**), and also how emissions from electricity are reported:

- This reporting rule relates to consumption that is 'unallocated'; this is the difference between the electricity that is supplied to Gibraltar's electricity network and the electricity that GibElec bill to customers.
- 'Unallocated' electricity is assumed to be lost through transmission and distribution and is therefore reported under scope 3.
- 'Unallocated' electricity is over double in 2017 than 2016 (but only 3% lower in 2017 than 2015r) meaning a smaller proportion is allocated to scope 2, rather than scope 3, hence a decrease in scope 2 electricity emissions.
- Combined scope 2 and 3 electricity emissions have decreased by 15% since 2016r and 20% since 2015r in line with electricity generation reported under Scope 1.

²⁵ not included in Gibraltar's 'manageable' emissions

Figure 5-6 - Gibraltar's emissions from electricity consumption/generation



The implied emission factor (IEF) for electricity considerably lower in 2017 than in 2016r; this means that less electricity was reported as generated per volume of fuel reported as consumed at the power stations in 2017 than 2016r, implying that the efficiency of electricity generation has increased. However, with only three years of data, it is not easy to draw firm conclusions. **Table 5-5** below shows a comparison of electricity data and IEF between years.

Emissions from the generation of electricity are calculated on the basis of total fuel consumed in the power stations. There have been no significant changes to the methodology and assumptions used: all years have estimated emissions using the NAEI emission factor for Gas Oil in power stations, and follow a simple approach of multiplying the fuel use (and activity data) by this emission factor.

At present the inventory calculation process is not sensitive enough to see a change in the IEF given different generation technologies, where they are using the same fuel. This is because more detailed information on plant generation characteristics is required to estimate the non-CO₂ gases. The CO₂ emissions remain unchanged as the quantity of carbon is fixed for combusting a given amount of fuel. In addition, because the supply of electricity from multiple sources is treated as a 'Gibraltar grid', the fuel and electricity outputs are aggregated to generate the IEF that represents an average across all generation technologies. Typically, the IEF will change as the balance of fuel and combustion types changes, for example a large input of renewables would increase the overall level of supply but without increasing the overall consumption of fuel, therefore the IEF would decrease. Likewise switching from gas oil to natural gas. Small changes between use of fuels in different plants is less likely to show a large impact. It is important to note however, that the IEF is only an indicative number that allows for the disaggregation of electricity emissions across end users based on estimated consumption.

Table 5-5 – Comparison of electricity consumption and production data, 2015r, 2016r and 2017

	2015r	2016r	2017	% change	
				2015r - 2017	2016r - 2017
Kt fuel consumed	60.0	56.9	48.2	-20%	-15%
GWh electricity produced	223.9	200.0	222.3	-1%	11%
Implied Emission Factor (IEF) (kt CO ₂ e/GWh)	0.9	0.9	0.7	-20%	-24%

Stationary Energy – Fuel combustion

Emissions from stationary fuel combustion have reduced by 4% and 6% between 2017 and 2015r and 2016r (respectively); this is due to slightly less diesel used for back-up electricity generation in the hospital and hotels. There are also small change in the Defra Conversion factors used for each year.

Road Transport

Road transport emission are only reported for Scope 1, with all fuel consumed by Gibraltarian vehicles reported in boundary²⁶. Whilst recalculations of emissions for each particular fuel type are more significantly affected by this, overall emissions from the road transport sector have reduced by only 1.0% and 0.2% in 2015r and 2016r respectively. Since emissions from this sector are largely dictated by fuel import statistics received from HM Customs, a slight increase in petrol imports (8%) causes a small increase in emissions from road transport in 2017.

It should also be noted that there have been significant recalculations to the fuel consumed in the road transport sector. After consultation with HM Customs in March 2019, assumptions made to interpret raw import statistics were revisited. Harmonisation of customs declarations with EU standards over 2017 and 2018 are thought to have improved the reliability of import statistics, particularly for the importation of road-worthy diesel, which is thought to have confused with other forms of diesel, such as marine diesel, and gas oil. As a result, estimates for imported fuel, and therefore the fuel consumed on Gibraltar's road network, have been revised for both 2016 and 2015.

Waste

Total reported emissions from Waste have seen a reduction of 13% and 10% in 2017 compared to 2015r and 2016r (respectively); this is due to decreases in the amount of waste that has been composted and sent to landfill. Emissions from the incineration of clinical waste have reduced slightly since 2015r.

Wastewater emissions from wastewater have increased slightly as a result of population growth, as the methodology is a Tier 1 population-based approach.

Waterborne navigation

Scope 1 emissions from waterborne (private boats) have changed significantly and are the most important recalculation in comparison to the 2018 submission. The most important factor in the recalculation is the revision to the assumptions about the fuel consumed by vessels included under Scope 1. Stakeholder discussions with the Gibraltar Port Authority in March 2019 revealed that the previous estimates of fuel use were overestimates. Previously, historic time-series analyses had suggested that diesel and petrol consumption amounted to about 30% and 10% of the imported fuel for a given year. However, following our discussions with the Gibraltar Port Authority, these assumptions were reduced to 15% and 5%. In future years, the inventory agency hopes to use more comprehensive fuel use data, such as statistics on the supply of automotive gas oil to vessels from refuelling activities which is now thought to exist in a usable format, but this assumption is kept at present as a conservative estimate of fuel use from this sector. In addition, these changes are influenced by the recalculations to total fuel imports described above. In total, emissions from this sector have reduced by 36% and 20% in 2015r and 2016r respectively a result of this recalculation, and account for the majority of the recalculations to the overall Gibraltar GHG inventories.

Scope 3 emissions have not been revised in either the 2015r or 2016r inventories.

Aviation

Emissions from aviation are around 36% higher in 2017 than 2015r, likely due to an increased number of flights to London Gatwick, London Heathrow and Manchester between 2015r and 2016r. Between 2016r and 2017, aviation emissions have increased by 4%, largely due to an increased number of flights to London Gatwick and Manchester.

Emissions from aviation are 3% lower 2017 compared to both the 2015r and 2016r inventories, as a result of revised assumptions on the fuel consumption at different stages of a flight stage, aligning with the NAEI.

²⁶ Consistent with GPC methodologies and best practice, where a robust method for splitting in-boundary and out of boundary emissions cannot be undertaken, fuel sales are reported under Scope 1. Sales to non-Gibraltarian vehicles is considered outside of Scope.

IPPU

There are still no Industrial Process emissions in Gibraltar, but Product Use emissions remain a significant source of scope 1 emissions (accounting for 13%). IPPU emissions have decreased by 4% in comparison to 2015r. This is a small decrease in terms of total tonnes of CO_{2e} and follows the UK trend for products including aerosols, firefighting, foams, refrigeration and air-conditioning.

Other Scope 3

Other Scope 3 includes emissions from out-of-scope shipping traffic (e.g. bunkering) and out-of-scope fuel use by non-Gibraltarian vehicles. These emissions have increased by 8% and 4% compared to the 2015r and 2016r inventories respectively. Emissions from other scope 3 sources show minor recalculations, a result of the changes in fuel import statistics discussed above influencing the road transport element of this sector.

More information on the specific reasons for changes between the 2017 and previous inventories are given in **Appendix 2**.

5.5 Reducing emissions in Gibraltar

An emissions inventory is an ongoing tool for understanding and reporting emissions, and allows the identification of major sources and priority areas for mitigation. From the results presented it can be seen that there are some areas where efforts should be focussed. Recommendations are therefore as follows:

- Stationary energy (in this instance nearly entirely electricity production) is the highest contributor to manageable emissions, and as such efforts should be focussed on both reducing consumption and decarbonising supply. A full assessment of both supply side mitigation options, and demand-side efficiency, should be undertaken.
- For transport, aviation and private marine are difficult to influence. However, scope 1 road transport, although relatively small compared to sources such as stationary energy, is significant given the small size of the territory. Road transport emissions would also be possible to influence through local policy measures. Gibraltar should therefore consider ways to reduce road transport, particularly car use.
- Proportionally, waste and IPPU emissions are aligned with the performance break-down in **Figure 5-4**, but in absolute terms, emissions from these sectors remain high. Gibraltar should therefore undertake research to identify methods to both reduce overall waste, increase recycling, and find alternative lower impact disposal methods.
- Individual industries that are high energy users should be identified and ways to reduce energy consumption investigated.

It is beyond the scope of this work to make detailed recommendations for mitigation measures beyond identifying the above major sources. However, it is recommended that should Gibraltar want to make significant improvements to its total emissions, a separate and detailed study of mitigation options and potential reductions in different sectors should be carried out.

5.6 Inventory accuracy

In compiling the Gibraltar City Inventory, there are a number of assumptions that have had to be made in the compilation and calculations that will have impacts on the accuracy of the data. The largest sources have been calculated with a high level of confidence, due to the presence of, for example, energy import statistics, detailed shipping records and the clearly bounded nature of activities (such as electricity generation). Some of the more minor assumptions relate to interpretation of data (such as units or fuel types where not consistent with the International System of Units (SI units), for instance). Most assumptions relate to methods of allocation within sectors, so the total inventory is associated with low uncertainty, but the sector allocations are more uncertain.

All assumptions have been documented in the relevant methodology section, but **Table 5-6** summarises some of the main assumptions and possible impacts on the data. A formal uncertainty analysis was not undertaken on the inventory as it was beyond the scope of this work.

Table 5-6 - Summary of assumptions and impact on inventory totals

Emission or data source	Assumption	Possible impact	Improvement
Proxy indicators	2015 values used as 2016/2017 values not available	It is unlikely that using 2015 values for proxy indicators (such as population) will have had a large impact on emission sources but it will be important to update to the correct year when available	<p>Latest year data for key indicators</p> <p>Consider working with the Abstract of Statistics team to obtain official correct year figures in advance of publication in future cycles</p>
Electricity allocation to end users	Electricity could only be allocated accurately for some users (domestic was based on tariffs and others were based on billings data) requiring allocation by proxy indicator	Ultimately, the total electricity emissions remain unchanged as this is an allocation issue. It is possible that some users have been over or under estimated and the emissions details possible for each end user is limited	<p>Billings data for other key sectors (such as public sector buildings, port, airport, retail) to allow better allocation</p> <p>It is anticipated that discussions and improvement work by GEA and AquaGib to refine and improve the tariffs and reporting by high users will improve the granularity of consumption data available in future years</p>
Transport activity data	Transport emissions were calculated by generating implied fuel consumption data based on the vehicle fleet. Actual information on vehicle movements was not available, so it was not possible to establish the proportion of travel in-boundary and out of boundary. It was therefore assumed that all fuel sold to Gibraltar vehicles was used in-boundary and all non-Gibraltar out of boundary	It is likely that the allocation of emissions has low accuracy. The implied vehicle kilometres (vkm) are for Gibraltar vehicles and are higher than would be expected for a region of this size. Therefore, it is likely this is an over estimate of in-boundary emissions. It is probable that some proportion of the Gibraltar fuel sales should be allocated to transboundary emissions, but it is not possible to distinguish. The lack of vehicle activity data also makes it difficult to account for off-road vehicles and public transport. The fuel import data provides the overall fuel balance, but in the transport sector some	<p>Data on household travel habits, in particular activity data to enable a better understanding of annual distance travelled by vehicle type (car/heavy goods vehicle (HGV)/light goods vehicle (LGV)/motorcycle, and private, commercial, public) would enable a better characterisation of vehicle emissions and improved allocation to end users.</p> <p>Understanding annual distance travelled by vehicle types can also be achieved through obtaining more detailed vehicle licensing data from the Department of Transport. By recording vehicle mileage during vehicle MOTs, high</p>

Emission or data source	Assumption	Possible impact	Improvement
		<p>of this is allocated to 'outside of scopes' as it is deemed to be 'exported' by non-Gibraltarian drivers. Therefore, the proportion of emissions from fuel imported that is allocated to Gibraltar is possibly over estimated. We also do not estimate the amount of fuel bought by Gibraltarians while outside of Gibraltar, which will lead to a small under estimate.</p> <p>Also, for 2017, we did not receive updated fleet data so assumptions have been made to update the fleet data.</p>	<p>quality data on the annual distance travelled by each vehicle will be available.</p> <p>It may also be possible to use ANPR technology, alongside the vehicle licensing information, to understand the split of vehicles travelling within and outside Gibraltar. This will give a far more accurate representation of the split of in- and out-of-boundary journeys than is currently available.</p>
Aviation activity data	Aviation was calculated on a bottom-up basis and was based on the number of flights, assumptions on the plane class and the expected distance flown. 'Unscheduled' flights were omitted as they were evidently linked to very small planes, for which we had limited emissions and fuel consumption estimates	There is some uncertainty on how much fuel would be used on journeys, the actual distance travelled and the validity of some of our assumptions. Additionally, the omission of the 'unscheduled' flights will lead to a small under estimate	Access to aviation fuel sales in Gibraltar would enable verification of bottom-up calculated fuel use data. This would reduce uncertainty as fuel sold gives a strong indication of the fuel use on outgoing journeys. This would also remove the possible under estimate due to the omission of unscheduled flights
Private marine emissions	A proportion of gas oil and petrol included in the fuel import data is used by private boats. This has been estimated at 15% and 5% of total demand for gas oil and petrol respectively in 2017, based on discussions with Gibraltar Port Authority and historic time-series analyses of fuel imports.	This assumption is supported by analysis of historical data trends. However, it is possible that this is not an accurate estimation of marine fuel use and is not sensitive to any trends in activity.	<p>Activity data, such as fuel sales at marine filling stations, would improve calculations of this emission source significantly.</p> <p>Information on the movements of private marine boats would also improve estimations from this sub-sector.</p>
Shipping activity data	Activity data for ships were estimated through calculations of distance travelled to and from other	It is possible that the ship classes and average fuel consumptions taken	The estimations in this sector are now based on a large amount of reliable and accurate

Emission or data source	Assumption	Possible impact	Improvement
	ports. This provided an indicator of fuel consumption per journey	form the EMEP/EEA air pollutant emission inventory guidebook 2016 do not accurately match the ships visiting Gibraltar. However, it is likely that any impact here is small	data. Accuracy could be further improved through use of ship specific fuel consumption and emission rates, technologies etc. to replace use of EMEP/EEA defaults.
Shipping activity data	Where it wasn't clear which port was being referred to in the port statistics, we have used a weighted, ship-type specific average distance to estimate more representative vessel journey lengths. Previously, simple average journey lengths were applied.	This increases the uncertainty in the distance travelled estimates and, hence, fuel consumption estimates	Complete, clear records of destination ports would reduce the need for assumptions to be applied
IPPU activity data	Little data existed on IPPU emissions for Gibraltar so these were estimated using UK data and proxy indicators (population, GDP) (with the exception of N ₂ O for anaesthesia and MDIs which have been accurately estimated).	It is possible that the Gibraltar case differs from the UK, particularly for air-conditioning units, which may be under estimated.	Latest year indicator data and Gibraltar-specific information on relevant product use, e.g. numbers of air-conditioning units, solvent use, etc. Some of this information is available (e.g. the number of refrigerators imported into Gibraltar); however, information on the current stock of such products in Gibraltar is not currently available.
Waste water emission calculation	Emissions were calculated using a default emission factor for wastewater to sea, lakes and rivers	It is likely that this has resulted in an over estimate of CH ₄ as sea water is less stagnant than lakes and inland waterways, so there will likely be less anaerobic decomposition	This is a very small inventory source. Improvements to estimates would require a level of work beyond the significance of the source
Waste composition data and disposal	The composition of municipal solid waste (MSW) arisings were estimated by applying the results of the 2015 Waste Characterisation Study to the total reported MSW. Therefore, this assumption is based on waste	It is probable that the fraction of waste recycled has been over estimated and emissions are, therefore, an under estimate. There are also assumptions about waste treatment in	Data from final processing rather than composition of waste exported should be obtained to improve this estimate. This is one of the inventory emission sources where local activity and decisions

Emission or data source	Assumption	Possible impact	Improvement
	<p>collection data, rather than final processing in Spain.</p> <p>In general, the waste sector has a lot of assumptions about composition and disposal/treatment methods</p>	<p>Spain which could result in uncertainty of the estimates. Overall, this is one of the smaller sources, so is less of a priority. However, it is possible that should the recycled fraction be lower, the source would have a greater overall emission share</p>	<p>can have a significant impact. Therefore, effort should be made to better characterise waste and understand disposal processes.</p>

6 Recommendations

This chapter sets out a series of recommendations following the compilation of this city-level greenhouse gas (GHG) inventory for Gibraltar. These recommendations include:

- Improving inventory compilation and future year reporting.
- Quality assurance, quality control (QAQC) and verification.

6.1 Improving inventory compilation and future year reporting

Inventories are very much intended to be a 'live' reporting tool and, as such, should be subject to regular revision and improvement. There are two particular aspects that should be considered by Gibraltar for ongoing improvement of compilation and future reporting of the inventory. These are:

- Improved activity data collection and management, including sectoral allocations.
- Recalculations and tracking emissions over time.

6.1.1 Improved activity data collection and management

All inventories have scope for improvement of data collection and management. The collection of data is often the most time consuming and challenging aspect of the inventory, so adequate time needs to be dedicated to this stage. The challenge is often that third parties hold the information that is required or that it is not available at all. It is vital to clearly express the data required – units, scope, boundaries, time period, sources and activities. It should also be requested that each data source is provided with an explanatory note and a contact for any queries. Failure to do so often results in incomplete data, the wrong data and a lack of transparency of how the data was compiled. An inventory is only ever as good as the data that underpins it. With this in mind, it is important to acknowledge that whilst data quality can be maximised, it can never be ensured. In addition, new data, improved information or clarity of assumptions may be developed over time, leading to recalculations and changes.

Under the Gibraltar inventory programme, a transparent and rigorous process of data requests, supply, processing and documentation has been implemented. Key to this has been the involvement of stakeholders and data suppliers, supporting the process of data identification, availability and transparency.

Currently and going forward, data required for Gibraltar's inventory is/will be requested during the summer. Data templates have been developed which are sent to data providers to encourage the provision of all the required data. The data templates provide space to enter the required data, as well as accompanying information (such as data quality, how the data was compiled, the period the data covers, etc) for QA/QC purposes. An example of a Gibraltar city inventory data template is given in **Appendix 5**. Data templates will be improved over time, working with data suppliers, to make the data collection process as efficient as possible.

Whilst the data templates aim to capture all relevant information, it should be acknowledged that these templates are not compulsory, and many suppliers will find it easier to provide data in other formats. Where this is the case, or supporting information is not clear or not provided, there are risks that data quality will be compromised. Efforts will therefore be made every year to engage data suppliers early and ensure that the principles of data quality can be maximised.

Currently, there are no formal agreements between the Government of Gibraltar and the data providers. Data supply agreements have been drafted, and are to be formally put in place, to ensure to consistent, timely and reliable supply of data for use in the inventory.

Appendix 4 sets out the minimum data requirements needed for estimation of emissions in each sector, and the data required for disaggregation and verification.

6.1.2 Recalculations and tracking emissions over time

It is important to track emissions over time to provide information on historical emissions trends, and the effects of policies and actions to reduce emissions at the city level.

As far as is possible, the time series should be calculated using the same methods, data sources and boundary definitions in all years to ensure consistency. Using different methods, data or applying different boundaries in a time series could introduce bias because the estimated emissions trend will reflect real changes in emissions or removals and the pattern of methodological refinements.

Significant changes may occur over time, which will alter the historical emissions profile, making meaningful comparisons over time difficult. To maintain consistency over time, historical emissions data from a base year inventory will have to be recalculated. This should also occur if methods change and data improve.

A number of recalculations have been made to the 2015 and 2016 inventories, based on data improvements realised through the 2017 inventory process; these are explained in **Appendix 3**.

The GPC sets out examples of the kinds of significant changes a city might experience that should trigger a recalculation. These are set out below and examples given in **Table 6-1**.

Structural changes in the assessment boundary. This may be triggered by adjustment in a city's administrative boundary, or changes in inclusion or exclusion of activities within the city boundary. For example, a category previously regarded as insignificant has grown to the point where it should be included in the inventory. However, no emissions recalculations are needed for activities that either did not exist in the base year, or reflect a natural increase or decrease in city activities ('organic growth').

- For Gibraltar, it will be necessary to track land use activities, particularly land reclamation as an extension of the territory and the emission implications of greater land area over time. The treatment of sources such as international bunkering, shipping and road vehicle fuel export may also be possible areas for change to assessment boundary (inclusion/exclusion) over time. Sources that have not been estimated should also be considered for inclusion.

Changes in calculation methodology or improvements in data accuracy. A city might report the same sources of GHG emissions as in previous years, but measure or calculate them differently. Changes resulting in significant emission differences should be considered as recalculation triggers, but any changes that reflect real changes in emissions do not trigger a recalculation. Sometimes, the more accurate data input may not reasonably be applied to all past years, or new data points may not be available for past years. The city may then have to back cast these data points, or the change in data source may simply be acknowledged without recalculation. This acknowledgement should be made in the report each year to enhance transparency. Otherwise, new users of the report in the two or three years after the change may make incorrect assumptions about the city's performance.

- There is a likelihood that Gibraltar activity data can be improved over time to give more accurate estimates. This is dealt with in more detail below and was discussed in **Section 5.4**.

Discovery of significant errors. A significant error or a number of cumulative errors that are collectively significant, should also be considered as a reason to recalculate emissions.

- QAQC procedures on this inventory should have minimised errors, but should an error have occurred, recalculations will be required.

Table 6-1: Examples of recalculation triggers, from Table 11.2 of GPC

Trigger	Example change	Recalculation needed	No recalculation needed
Changes in assessment boundary	A community is included in or set aside from a city's administrative boundary	X	
	Change in goal boundary from BASIC to BASIC+, or from six GHGs to seven GHGs	X	
	Shut down of a power plant		X
	Build of a new cement factory		X
Changes in calculation methodology or improvements in data accuracy	Change in calculation methodology for landfilled municipal solid waste (MSW) from <i>Mass Balance Method</i> to the <i>First Order Decay Method</i>	X	
	Adoption of more accurate local emission factors, instead of a national average for scope 2 emissions	X	
	Change in electricity emission factor due to energy efficiency improvement and growth of renewable energy utilisation		X
Discovery of significant errors	Discovery of mistake in unit conversion in formula used	X	

Whether recalculation is needed depends on the significance of the changes. The determination of a significant change may require taking into account the cumulative effect on base year emissions of a number of small changes.

6.2 QAQC and verification

6.2.1 Quality control

Quality control (QC) is a set of technical activities that measure and control the quality of the inventory as it is being developed. They are designed to:

- Provide routine and consistent checks to ensure data integrity, correctness and completeness.
- Identify and address errors and omissions.
- Document and archive inventory material and record all QC activities.

QC activities include accuracy checks on data acquisition and calculations, and the use of approved standardised procedures for emission calculations, measurements, estimating uncertainties, archiving information and reporting. Higher tier QC activities include technical reviews of source categories, activity and emission factor data, and methods.

A number of QC checks were undertaken in the compilation of the inventory; these included:

- Mass balance checks – fuel data 'used' versus fuel data 'supplied' for Gibraltar should balance.
- Implied Emission Factors (IEFs) – checks against UK GHG inventory to ensure the order of magnitude is what would be expected.
- Timeseries checks – checks against previous year to assess data accuracy and completeness.
- Spreadsheet functions – manual checks that formulae are working as expected.
- Consistent labelling, file revisions (e.g. dated file extensions).
- Documentation on spreadsheets, with details of calculation method, assumptions, emission factors and data quality.

6.2.2 Quality assurance

Quality assurance (QA) activities include a planned system of review procedures conducted by personnel not directly involved in the inventory compilation/development process. Reviews, preferably performed by independent third parties, should take place when an inventory is finalised following the implementation of QC procedures. Reviews verify that data quality objectives were met and that the inventory represents the best possible estimates of emissions – and sinks given the current state of scientific knowledge and data available.

Several QA reviews were undertaken by internal inventory experts for the calculations for each sector and of methodologies used across the inventory.

Table 6-2: Quality Checking

Sector	Reviewer(s)
Stationary Energy	Luke Jones
Waste	Luke Jones
Road transport	Peter Brown
Aviation	Luke Jones
Shipping	Luke Jones and Ellie Kilroy
IPPU	Luke Jones

In addition, quality checks of the final reported data to ensure consistency with the GPC and complete and transparent reporting of the final results, and documentation of methods and results in this report are also carried out by the Knowledge Leadership and project management team.

6.2.3 Verification

Verification can be used to increase credibility of publicly reported emissions information with external audiences and increase confidence in the data used to develop climate action plans, set GHG targets and track progress.

Verification involves an assessment of the completeness, accuracy and reliability of reported data. It seeks to determine if there are any material discrepancies between reported data and data generated from the proper application of the relevant standards and methodologies. It does this by making sure that the reporting requirements have been met, that the estimates are correct and that the data sourced are reliable.

No verification was carried out on this report or the underlying data, due a lack of defined verification processes and bodies to carry this out.

Appendices

Appendix 1: Common Reporting Framework (CRF) for 2017

Appendix 2: Detailed reasons for changes between 2017 and previous inventories

Appendix 3: Recalculations

Appendix 4: Data recommendations

Appendix 5: Data collection template

Appendix 1 – Common Reporting Framework (CRF) for 2017

GHG Emissions Source (By Sector and Sub-sector)	Total GHGs (metric tonnes CO ₂ e)				Comments
	Direct	Other Direct	Indirect	Total	
STATIONARY ENERGY					
Residential buildings	NO	9,085	43,348	52,433	
Commercial buildings and facilities	1,423	17,075	81,477	99,976	
Institutional buildings and facilities	NO	113	538	651	Reported under '1.6 Non-specified sources' in the GPC
Industrial buildings and facilities	NO	IE	2,681	2,681	Reported under '1.4 Energy industries' in the GPC
Agriculture	NO	NO	NO		
Fugitive emissions	NO	NO	NO		
SUB-TOTAL	1,423	26,273	128,044	155,740	
TRANSPORTATION					
On-road	55,438	IE	NO	55,438	
Rail	NO	NO	NO		
Waterborne navigation	12,576	262,783	NO	275,359	
Aviation	NO	38,482	NO	38,482	
Off-road	IE	IE	NO		
SUB-TOTAL	68,014	301,265		369,279	
WASTE					
Solid waste disposal	NO	15,782	NO	15,782	
Biological treatment	NO	1,489	NO	1,489	
Incineration and open burning	NO	398	NO	398	
Wastewater treatment and discharge	NO	1,790	NO	1,790	
SUB-TOTAL		19,460		19,460	
INDUSTRIAL PROCESSES and PRODUCT USES					
Industrial Process	NO	NO	NO		
Product Use	NO	10,591	NO	10,591	
SUB-TOTAL		10,591		10,591	
AGRICULTURE, FORESTRY and OTHER LAND USE					
Livestock	NO	NO	NO		
Land use	NO	NO	NO		
Other AFOLU	NO	NO	NO		
SUB-TOTAL					
TOTAL	69,437	357,588	128,044	555,069	
ENERGY GENERATION					
Electricity-only generation	154,317			154,317	
CHP generation	NO				
Heat/cold generation	NO				
Local renewable generation	NE				
SUB-TOTAL	154,317			154,317	

Appendix 2 – Detailed reasons for changes between 2017 and previous inventories

This table provides a summary of the reasons for changes in emissions in sub-sectors. Sub-sectors not included in this table did not show any significant change in emissions between years.

Source	Change between current year (2017) base year (2015r)	Change between current year (2017) and previous year (2016r)	Reason
Stationary Energy			
Electricity generation	Decrease	Decrease	Emissions from electricity generation have decreased by 19% since 2015r and 15% since 2016r; this is due to less fuel being used to generate a unit of electricity, implying improvements in efficiency at Gibraltar's electricity power stations.
Scope 2 Electricity	Decrease	Decrease	<p>Scope 2 electricity emissions (electricity consumption) have decreased by over 30kt CO_{2e}, or 19%, since 2015r. This is due to less fuel being used to generate electricity, rather than electricity consumption itself reducing.</p> <p>In 2016r, more electricity was assigned to scope 2 electricity consumption compared to scope 3 transmission and distribution losses (as more electricity was 'billed' of the total supplied to the grid), which led an increase in overall scope 2 electricity emissions in 2016r. Less electricity was therefore considered as scope 3 transmission and distribution losses, or unbilled electricity in 2016r compared to 2015r and 2017.</p> <p>Since 2016, residential and commercial electricity use has increased, but electricity use by the 'power' tariff and by the MOD have decreased.</p>
Scope 3 Electricity	Decrease	Increase	The difference between the amount of electricity supplied to the Gibraltar electricity network and the amount of electricity that is billed for by AquaGib (reported under scope 2) is assumed to be the transmission and distribution losses across the network. In 2016, there was around half the amount of this 'unallocated' electricity reported than 2015 and 2017; this could be due to improvements in the way electricity is billed to consumers (with subsequent increases in scope 2, see above row) or due to actions to improve electricity losses across the network. The 2017 results are more consistent with the 2015 results, hence an increase between 2016r and 2017.

Transportation			
Scope 1 On-road transportation	Increase	Increase	Trends in road transport emissions are dictated principally by changes the fuel imported into Gibraltar. For example, an 8% increase in petrol imports is reflected in the emissions statistics from this particular fuel. To a lesser extent, changes in fleet composition, with greater penetration of Euro 6 vehicles which tend to be more fuel efficient and have differing methane and nitrous oxide factors cause emission trends, but these are much less important.
Scope 1 Waterborne navigation	Increase	Increase	Trends in Scope 1 waterborne emissions are dictated principally by changes the fuel imported into Gibraltar. For example, an 8% increase in petrol imports is reflected in the emissions statistics from this particular fuel.
Aviation	Increase	Increase	Emissions from aviation are around 36% higher in 2017 than 2015r, likely due to an increased number of flights to London Gatwick, London Heathrow and Manchester between 2015r and 2016r. Between 2016r and 2017, aviation emissions have increased by 4%, largely due to an increased number of flights to London Gatwick and Manchester.
Waste			
Landfill and Biological treatment of waste	Decrease	Decrease	Emissions from Waste are around 13% and 10% lower in 2017 than 2015r and 2016r (respectively); this is due to decreases in the amount of waste that has been composted and sent to landfill.
Incineration of waste	Increase	Increase	Scope 1 emissions from waste are attributable to the incineration of clinical waste within Gibraltar. Emissions from this activity have increased since 2015r due to an increase in clinical waste arisings treated by incineration. Scope 3 emissions from incinerating waste outside of Gibraltar have decreased compare to 2016, but increase compared to 2015.
Wastewater	Increase	Increase	Wastewater emissions have increased along with population growth.
IPPU			
Product use	Decrease	Decrease	There are still no Industrial Process emissions in Gibraltar, but Product Use emissions remain a significant source of scope 1 emissions (accounting for 13%). IPPU emissions have decreased by 4% in comparison to 2015r. This is a small decrease in terms of total tonnes of CO ₂ e and follows the UK trend for products including aerosols, firefighting, foams, refrigeration and air-conditioning.
Other Scope 3			

Road Transport	Increase	Decrease	Revisions to assumptions made during cleaning and handling of raw fuel import statistics are responsible for small declines in the use of petrol and diesel on-road, both by non-Gibraltarian vehicles and Gibraltarian vehicles (the latter as a part of Scope 1)
----------------	----------	----------	--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Table A-1 presents the magnitude of these changes in metric tonnes CO₂e and % between the 2015r and 2017 inventories. **Table A-2** presents the magnitude of these changes in metric tonnes CO₂e and % between the 2016r and 2017 inventories.

Note that negative numbers reflect a reported reduction and positive numbers a reported increase.

Table A-1: comparison of changes between 2015r-2017 by sector (excluding those not occurring)

GPC ref No.	GHG Emissions Source (By Sector and Sub-sector)	Change 2015r vs. 2017 (metric tonnes CO2e)				Change (%)			
		Scope 1	Scope 2	Scope 3	Total	Scope 1	Scope 2	Scope 3	Total
I	STATIONARY ENERGY								
I.1	Residential buildings	NO	-8,605	-2,114	-10,720	NO	-17%	-19%	-17%
I.2	Commercial and institutional buildings and facilities	-64	-20,675	-4,944	-25,683	-4%	-20%	-22%	-20%
I.4.1/2/3	Energy industries	NO	-284	IE	-284	NO	-10%	IE	-10%
I.4.4	Energy generation supplied to the grid	-37,635				-20%			
I.6	Non-specified sources	NO	-831	-182	-1,013	NO	-61%	-62%	-61%
SUB-TOTAL	(city induced framework only)	-64	-30,395	-7,240	-37,700	-4%	-19%	-22%	-19%
II	TRANSPORTATION								
II.1	On-road transportation	2,602	NO	NO	2,602	5%	NO	NO	5%
II.3	Waterborne navigation	190	NO	72,884	73,073	1.5%	NO	38%	36%
II.4	Aviation	NO	NO	10,115	10,115	NO	NO	36%	36%
SUB-TOTAL	(city induced framework only)	2,792	NO	82,999	85,791	4%	NO	38%	30%
III	WASTE								
III.1.1/2	Solid waste generated in the city	NO		-2,591	-2,591	NO		-14%	-14%
III.2.1/2	Biological waste generated in the city	NO		-244	-244	NO		-14%	-14%
III.3.1/2	Incinerated and burned waste generated in the city	72		-70	2	26%		-57%	1%
III.4.1/2	Wastewater generated in the city	NO		44	44	NO		3%	3%
SUB-TOTAL	(city induced framework only)	72		-2,861	-2,789	26%		-13%	-13%
IV	INDUSTRIAL PROCESSES and PRODUCT USES								
IV.2	Emissions from product use occurring within the city boundary	-470			-470	-4%			-4%
SUB-TOTAL	(city induced framework only)	-470			-470	-4%			-4%
VI	OTHER SCOPE 3								
VI.1	Other Scope 3			247,219	247,219			8%	8%
TOTAL	(city induced framework only)	2,329	-30,395	320,116	292,050	3%	-19%	10%	8%

Table A-2: comparison of changes between 2016r-2017 by sector (excluding those not occurring)

GPC ref No.	GHG Emissions Source (By Sector and Sub-sector)	Change 2016r vs. 2017 (metric tonnes CO ₂ e)				Change (%)			
		Scope 1	Scope 2	Scope 3	Total	Scope 1	Scope 2	Scope 3	Total
I	STATIONARY ENERGY								
I.1	Residential buildings	NO	-11,617	3,382	-8,235	NO	-21%	59%	-14%
I.2	Commercial and institutional buildings and facilities	-88	-25,498	5,977	-19,609	-6%	-24%	54%	-16%
I.4.1/2/3	Energy industries	NO	394	IE	394	NO	17%	IE	17%
I.4.4	Energy generation supplied to the grid	-27,879				-15%			
I.6	Non-specified sources	NO	-520	3	-517	NO	-49%	3%	-44%
SUB-TOTAL	(city induced framework only)	-88	-37,241	9,361	-27,967	-6%	-23%	55%	-15%
II	TRANSPORTATION								
II.1	On-road transportation	1,695	NO		1,695	3%	NO	NO	3%
II.3	Waterborne navigation	126	NO	-10,500	-10,374	1.0%	NO	-4%	-4%
II.4	Aviation	NO	NO	1,640	1,640	NO	NO	4%	4%
SUB-TOTAL	(city induced framework only)	1,821	NO	-8,860	-7,039	3%	NO	-3%	-2%
III	WASTE								
III.1.1/2	Solid waste generated in the city			-1,905	-1,905	NO		-11%	-11%
III.2.1/2	Biological waste generated in the city			-180	-180	NO		-11%	-11%
III.3.1/2	Incinerated and burned waste generated in the city	43		-82	-39	14%		-60%	-9%
III.4.1/2	Wastewater generated in the city			23	23	NO		1%	1%
SUB-TOTAL	(city induced framework only)	43		-2,144	-2,101	14%		-10%	-10%
IV	INDUSTRIAL PROCESSES and PRODUCT USES								
IV.2	Emissions from product use occurring within the city boundary	-284			-284	-3%			-3%
SUB-TOTAL	(city induced framework only)	-284			-284	-3%			-3%
VI	OTHER SCOPE 3								
VI.1	Other Scope 3			117,659	117,659			4%	4%
TOTAL	(city induced framework only)	1,492	-37,241	116,016	80,268	2%	-23%	3%	2%

Appendix 3 – Recalculations

This appendix covers the main recalculations between the 2018 submission and the 2019 revised inventories for 2015 and 2016. Recalculations with a very small or insignificant impact have not been covered.

Sector	Sector/sub-sector	Change in tonnes of CO ₂ e to 2015r	Change in tonnes of CO ₂ e to 2016r	Reason
II. Transport	II.1 - Road Transport	-523	-134	Revisions to assumptions made during cleaning and handling of raw fuel import statistics are responsible for small declines in the use of petrol and diesel on-road, both by Gibraltarian vehicles and non-Gibraltarian vehicles (the latter as a part of Other Scope 3)
II. Transport	II.3 Waterborne Navigation	- 28,604	- 28,541	Revisions to assumptions dictating the amount of fuel used by marina vessels after discussions with the Gibraltar Port Authority (now equivalent to 15% and 5% of diesel and petrol imports respectively), and assumptions made during cleaning and handling of raw fuel import statistics are responsible for large recalculations from this sector. These constitute the most important revisions across the whole city GHG inventory.

Sector	Sector/sub-sector	Change in tonnes of CO2e to 2015r	Change in tonnes of CO2e to 2016r	Reason
II. Transport	II.4 Aviation	-1,244	-954	These recalculations are as a result of revised assumptions on the fuel consumption at different stages of a flight (mainly taxing times), aligning with the NAEI.
III. Waste	III.1 Landfills	54	46	Total waste volumes include mattresses in the recalculated inventories
III. Waste	III.4 Wastewater	-175	-178	An assumption in the wastewater calculations has been amended. It was previously assumed that Gibraltar had garbage disposal, which refers to a unit installed under a kitchen sink that traps and shreds food waste into pieces small enough to pass through plumbing. It is now assumed that this is not the case.
IV. IPPU	IV.2 Product Use	-45	165	Small recalculations as a result of revisions to the UK F-Gas model which the Gibraltar inventory uses as a proxy for Product Use emissions.
VI. Other Scope 3	VI. Other Scope 3	-5,196	298	See II.1 Road transport

Appendix 4 – Data recommendations

The table below sets out the data requirements for each of the main sectors. It shows the minimum top-level data required for emission calculation, and the data required to enable a disaggregation of the data by end user and/or category. The Data for Verification column shows the data required to cross-check and verify the disaggregation of data.

Cells in grey indicate data that was not available for the Gibraltar 2016 inventory.

Sector	Minimum top level data	Data for disaggregation	Data for verification
Power	Fuel consumption for power (electricity) generation by fuel type <ul style="list-style-type: none"> - Gibraltar power station - 	Electricity produced in Gibraltar (total) Electricity consumed by sector (e.g. residential, commercial, Government/public services, Industrial) <ul style="list-style-type: none"> - Billings by tariff or end-user - Meter readings 	Total power (electricity) generation
Fuels/ combustion	Total fuel consumption by fuel type (non-electricity generation)	Fuel combustion locations End user sales / permits Total use by purpose (cooking stove, boiler etc.)	Not applicable
Transport (road)	Fuel import data by fuel type	Gibraltar vehicle licencing statistics End-user activity split : fuel use by vehicle type and purpose (including in and out of boundary – crossings of the Frontier) Fuel sold	Vehicle kilometre (vkm) data, by vehicle type and purpose
Marine (private)	Fuel import data	Fuel sold Fleet composition Fuel usage by marine use (boat type)	Not applicable
Shipping	Port activity <ul style="list-style-type: none"> - Number of ships - Types - Distance (origin/destination) 	Ship details (each) <ul style="list-style-type: none"> - Purpose - Class - Tonnage Purpose for calling (bunkers/non-bunkers)	Fuel sold
Off-road	Fuel sold	Licencing statistics for off-road fleet Fleet composition Fuel use by vehicle type	Vehicle kilometre (vkm) data or hours of use

Aviation (from CAA)	Numbers of flights and destinations Distances flown (origin/destination)	Fleet data (aircraft types)	Fuel sold
Waste	Total tonnage of waste Disposal methods	Tonnes / type - Biological content - Further information on the waste treatment process in Spain	Not applicable
Wastewater	Total volume of wastewater Biological content Treatment streams	Wastewater volume by sector	Population Average wastewater and biological content per person
Industrial Processes and Product Use	Numbers of products by type (e.g. A/C units, refrigerators, vehicle A/C) (some data available) Volumes of N ₂ O (hospital)	Numbers of products by end use sector	Population GDP Average product use / number per person / GDP

Appendix 5 – Data collection template

The below is an extract from the data collection template that has been sent to data providers during the data collection process.

Data

Please use this template as a guide to the information required for the GibEMIT programme

Please do not feel constrained by the template - tables can be altered or replaced as best suits the information available - but the information requested below should all be provided if possible

Specify sector relevant to this data
 Sector

Specify sub-sector(s) relevant to this data. Select additional sub-sectors in the next column if necessary
 Sub-Sector(s)

Note: Data should correspond to the 2015 calendar year and the boundary of the Government of Gibraltar

GPC Ref	Activity	Description	Amount	Units	Time-period	Data Quality	Description of the data/methods	Source
Compiler input	State the activity, site, end-user, fuel type, transport mode etc. e.g. "Residential Electricity consumption" "Buses"	If required, please provide further description of the data e.g. "total consumed electricity from all billings to residential electricity tariffs within Gibraltar" "Total diesel consumption in buses on route 1"	State the data value e.g. if 10 litres, state 10	Specify the unit, e.g. litres, km, kWh	Specify whether the data is annual (and the year), monthly (and the month) etc.	If possible, please give an indicative assessment of the quality of the data (L,M,H) Note - H is directly measured/recorded locally specific data, L is international default or highly modelled	If relevant, please give provide a brief description of any methods used to derive the data	If relevant, please state the source of this data, place of publication etc.

As well as collecting the actual activity data, additional information is also requested for quality control purposes; this information is presented below.

QC information required	Description of information required
Compiler	<i>Who compiled this data?</i>
Date created	<i>When was this data created/compiled?</i>
Source of data	<i>Where has this data come from?</i>
Data provided to	<i>Who has this data been provided for?</i>
Data purpose	<i>What has this data been provided for? Does this affect its use?</i>
Quality / Checking	<i>Has this data been checked by anyone? How has it been checked? Can you give an indication of the data quality?</i>
Data range / scope	<i>Time (e.g. date range) Geographic scope Installations/activities</i>
Notes/disclaimers	<i>Any other important information that the data recipient should be aware of? Are there missing years? Is this an estimate? Is this confidential?</i>



Ricardo
Energy & Environment

The Gemini Building
Fermi Avenue
Harwell
Didcot
Oxfordshire
OX11 0QR
United Kingdom

t: +44 (0)1235 753000
e: enquiry@ricardo.com

ee.ricardo.com